



2/2



Rec'd PCT/PTO 24 JAN 2005  
PCT/GB 3 / 0 0 3 1 7 1



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 21 AUG 2003

WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated

*P. McHoney*  
12 August 2003

Best Available Copy

24. DEC. 2002 12:14

GLOBAL IP PATENTS

NO. 2941 P. 2

Patents Form 1/77

Patents Act 1977  
(Rule 16)

THE PATENT OFFICE

24 DEC 2002

RECEIVED BY FAX

The  
Patent  
Office31DEC02 1774206-1 002734  
P01/7700 0.80-0230318.8**Request for grant of a patent***(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)*

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference

100804-2

2. Patent application number

*(The Patent Office will fill in this part)*

0230318.8

3. Full name, address and postcode of the or of each applicant *(underline all surnames)*AstraZeneca AB  
S-151 85 Sodertalje  
SwedenPatents ADP number *(if you know it)*

7822448004

If the applicant is a corporate body, give the country/state of its incorporation

Sweden

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent *(if you have one)*

Lucy Clare Padget

*"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)*AstraZeneca UK Limited  
Global Intellectual Property  
Mereside, Alderley Park  
Macclesfield  
Cheshire SK10 4TGPatents ADP number *(if you know it)*

8346762001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and *(if you know it)* the or each application number

Country

Priority application number  
*(if you know it)*Date of filing  
*(day / month / year)*

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
*(day / month / year)*8. Is a statement of inventorship and of right to grant of a patent required in support of this request? *(Answer 'Yes' if:*a) *any applicant named in part 3 is not an inventor, or*b) *there is an inventor who is not named as an applicant, or*c) *any named applicant is a corporate body.**See note (2))*

Patents Form 1/77

0056436 24-Dec-02 11:46

**Patents Form 1/77**

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 104

Claim(s) 3

Abstract 1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents  
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

Authorised Signatory

24/12/2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Jennifer C Bennett - 01625 230148

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

100804-2

- 1 -

**CHEMICAL COMPOUNDS**

This invention relates to chemical compounds, or pharmaceutically acceptable salts thereof. These compounds possess human 11- $\beta$ -hydroxysteroid dehydrogenase type 1 enzyme (11 $\beta$ HSD1) inhibitory activity and accordingly have value in the treatment of disease states including metabolic syndrome and are useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said compounds, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit 11 $\beta$ HSD1 in a warm-blooded animal, such as man.

10        Glucocorticoids (cortisol in man, corticosterone in rodents) are counter regulatory hormones i.e. they oppose the actions of insulin (Dallman MF, Strack AM, Akana SF et al. 1993; Front Neuroendocrinol 14, 303-347). They regulate the expression of hepatic enzymes involved in gluconeogenesis and increase substrate supply by releasing glycerol from adipose tissue (increased lipolysis) and amino acids from muscle (decreased protein synthesis and  
15    increased protein degradation). Glucocorticoids are also important in the differentiation of pre-adipocytes into mature adipocytes which are able to store triglycerides (Bujalska II et al. 1999; Endocrinology 140, 3188-3196). This may be critical in disease states where glucocorticoids induced by "stress" are associated with central obesity which itself is a strong risk factor for type 2 diabetes, hypertension and cardiovascular disease (Bjorntorp P &  
20    Rosmond R 2000; Int. J. Obesity 24, S80-S85)

It is now well established that glucocorticoid activity is controlled not simply by secretion of cortisol but also at the tissue level by intracellular interconversion of active cortisol and inactive cortisone by the 11-beta hydroxysteroid dehydrogenases, 11 $\beta$ HSD1 (which activates cortisone) and 11 $\beta$ HSD2 (which inactivates cortisol) (Sandeep TC & Walker  
25    BR 2001 Trends in Endocrinol & Metab. 12, 446-453). That this mechanism may be important in man was initially shown using carbenoxolone (an anti-ulcer drug which inhibits both 11 $\beta$ HSD1 and 2) treatment which (Walker BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159) leads to increased insulin sensitivity indicating that 11 $\beta$ HSD1 may well be regulating the effects of insulin by decreasing tissue levels of active glucocorticoids (Walker  
30    BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159).

Clinically, Cushing's syndrome is associated with cortisol excess which in turn is associated with glucose intolerance, central obesity (caused by stimulation of pre-adipocyte differentiation in this depot), dyslipidaemia and hypertension. Cushing's syndrome shows a



100804-2

- 2 -

number of clear parallels with metabolic syndrome. Even though the metabolic syndrome is not generally associated with excess circulating cortisol levels (Jessop DS et al. 2001; J. Clin. Endocrinol. Metab. 86, 4109-4114) abnormally high 11 $\beta$ HSD1 activity within tissues would be expected to have the same effect. In obese men it was shown that despite having similar or  
5 lower plasma cortisol levels than lean controls, 11 $\beta$ HSD1 activity in subcutaneous fat was greatly enhanced (Rask E et al. 2001; J. Clin. Endocrinol. Metab. 1418-1421). Furthermore, the central fat, associated with the metabolic syndrome expresses much higher levels of 11 $\beta$ HSD1 activity than subcutaneous fat (Bujalska J et al. 1997; Lancet 349, 1210-1213). Thus there appears to be a link between glucocorticoids, 11 $\beta$ HSD1 and the metabolic  
10 syndrome.

11 $\beta$ HSD1 knock-out mice show attenuated glucocorticoid-induced activation of gluconeogenic enzymes in response to fasting and lower plasma glucose levels in response to stress or obesity (Kotelevtsev Y et al. 1997; Proc. Natl. Acad. Sci USA 94, 14924-14929) indicating the utility of inhibition of 11 $\beta$ HSD1 in lowering of plasma glucose and hepatic  
15 glucose output in type 2 diabetes. Furthermore, these mice express an anti-atherogenic lipoprotein profile, having low triglycerides, increased HDL cholesterol and increased apolipoprotein AI levels. (Morton NM et al. 2001; J. Biol. Chem. 276, 41293-41300). This phenotype is due to an increased hepatic expression of enzymes of fat catabolism and PPAR $\alpha$ . Again this indicates the utility of 11 $\beta$ HSD1 inhibition in treatment of the  
20 dyslipidaemia of the metabolic syndrome.

The most convincing demonstration of a link between the metabolic syndrome and 11 $\beta$ HSD1 comes from recent studies of transgenic mice over-expressing 11 $\beta$ HSD1 (Masuzaki H et al. 2001; Science 294, 2166-2170). When expressed under the control of an adipose specific promoter, 11 $\beta$ HSD1 transgenic mice have high adipose levels of  
25 corticosterone, central obesity, insulin resistant diabetes, hyperlipidaemia and hyperphagia. Most importantly, the increased levels of 11 $\beta$ HSD1 activity in the fat of these mice are similar to those seen in obese subjects. Hepatic 11 $\beta$ HSD1 activity and plasma corticosterone levels were normal, however, hepatic portal vein levels of corticosterone were increased 3 fold and it is thought that this is the cause of the metabolic effects in liver.

30 Overall it is now clear that the complete metabolic syndrome can be mimicked in mice simply by overexpressing 11 $\beta$ HSD1 in fat alone at levels similar to those in obese man.

100804-2

- 3 -

11 $\beta$ HSD1 tissue distribution is widespread and overlapping with that of the glucocorticoid receptor. Thus, 11 $\beta$ HSD1 inhibition could potentially oppose the effects of glucocorticoids in a number of physiological/pathological roles. 11 $\beta$ HSD1 is present in human skeletal muscle and glucocorticoid opposition to the anabolic effects of insulin on protein turnover and glucose metabolism are well documented (Whorwood CB et al. 2001; J. Clin. Endocrinol. Metab. 86, 2296-2308). Skeletal muscle must therefore be an important target for 11 $\beta$ HSD1 based therapy.

Glucocorticoids also decrease insulin secretion and this could exacerbate the effects of glucocorticoid induced insulin resistance. Pancreatic islets express 11 $\beta$ HSD1 and carbenoxolone can inhibit the effects of 11-dehydrocorticosterone on insulin release (Davami B et al. 2000; J. Biol. Chem. 275, 34841-34844). Thus in treatment of diabetes 11 $\beta$ HSD1 inhibitors may not only act at the tissue level on insulin resistance but also increase insulin secretion itself.

Skeletal development and bone function is also regulated by glucocorticoid action. 11 $\beta$ HSD1 is present in human bone osteoclasts and osteoblasts and treatment of healthy volunteers with carbenoxolone showed a decrease in bone resorption markers with no change in bone formation markers (Cooper MS et al 2000; Bone 27, 375-381). Inhibition of 11 $\beta$ HSD1 activity in bone could be used as a protective mechanism in treatment of osteoporosis.

Glucocorticoids may also be involved in diseases of the eye such as glaucoma. 11 $\beta$ HSD1 has been shown to affect intraocular pressure in man and inhibition of 11 $\beta$ HSD1 may be expected to alleviate the increased intraocular pressure associated with glaucoma (Rauz S et al. 2001; Investigative Ophthalmology & Visual Science 42, 2037-2042).

There appears to be a convincing link between 11 $\beta$ HSD1 and the metabolic syndrome both in rodents and in humans. Evidence suggests that a drug which specifically inhibits 11 $\beta$ HSD1 in type 2 obese diabetic patients will lower blood glucose by reducing hepatic gluconeogenesis, reduce central obesity, improve the atherogenic lipoprotein phenotype, lower blood pressure and reduce insulin resistance. Insulin effects in muscle will be enhanced and insulin secretion from the beta cells of the islet may also be increased.

Currently there are two main recognised definitions of metabolic syndrome.

1) The Adult Treatment Panel (ATP III 2001 JMA) definition of metabolic syndrome indicates that it is present if the patient has three or more of the following symptoms:

➤ Waist measuring at least 40 inches (102 cm) for men, 35 inches (88 cm) for women;

100804-2

- 4 -


- Serum triglyceride levels of at least 150 mg/dl (1.69 mmol/l);
- HDL cholesterol levels of less than 40 mg/dl (1.04 mmol/l) in men, less than 50 mg/dl (1.29 mmol/l) in women;
- Blood pressure of at least 135/80 mm Hg; and / or
- 5 ➤ Blood sugar (serum glucose) of at least 110 mg/dl (6.1 mmol/l).

2) The WHO consultation has recommended the following definition which does not imply causal relationships and is suggested as a working definition to be improved upon in due course:

- The patient has at least one of the following conditions: glucose intolerance, impaired glucose tolerance (IGT) or diabetes mellitus and/or insulin resistance; together with two or more of the following:
- 10 ➤ Raised Arterial Pressure;
- Raised plasma triglycerides
- Central Obesity
- 15 ➤ Microalbuminuria

We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt thereof, are effective 11 $\beta$ HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndrome.

Accordingly there is provided the use of a compound of formula (I):



(I)

wherein:

Ring A is selected from aryl or heteroaryl;

- R<sup>1</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkanoyl, C<sub>1-6</sub>alkanoyloxy, N-(C<sub>1-6</sub>alkyl)amino, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, C<sub>1-6</sub>alkanoylamino, N-(C<sub>1-6</sub>alkyl)carbamoyl, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-6</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-6</sub>alkoxycarbonyl, N-(C<sub>1-6</sub>alkyl)sulphamoyl, N,N-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-6</sub>alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-6</sub>alkylene-Y- and heterocyclylC<sub>0-6</sub>alkylene-Y-; or two R<sup>1</sup> on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy group; wherein R<sup>1</sup> may be optionally substituted
- 25
- 30

100804-2

- 5 -

on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;

$n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

- $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkylS(O) <sub>$a$</sub> , wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl, carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}$ alkyl and heterocyclyl $C_{1-4}$ alkyl; or  $R^2$  and  $R^3$  together form oxo; wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{10}$ ;

$X$  is  $-CR^{11}R^{12}$ -,  $-S(O)_a$ -,  $-O$ -,  $-NR^{13}$ -,  $-C(O)$ -,  $-C(O)NR^{14}$ -,  $-NR^{15}C(O)$ -,  $-SO_2NR^{16}$ - or  $-NR^{16}SO_2$ -; wherein  $a$  is 0 to 2;

$r$  is 1 or 2;

$q$  is 0 or 1;

- 15  $p$  is 0 or 1;

Ring  $B$  is carbocyclyl or heterocyclyl; wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

- $R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O) <sub>$a$</sub> , wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}$ alkylene-Y- and heterocyclyl $C_{0-4}$ alkylene-Y-; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

$m$  is 0-3; wherein the values of  $R^6$  may be the same or different;

- $Y$  is  $-S(O)_a$ -,  $-O$ -,  $-NR^{20}$ -,  $-C(O)$ -,  $-C(O)NR^{21}$ -,  $-NR^{22}C(O)$ - or  $-SO_2NR^{23}$ -; wherein  $a$  is 30 0 to 2;

$R^7$ ,  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,

100804-2

- 6 -

*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein R<sup>7</sup>, R<sup>9</sup> and R<sup>18</sup> may be independently optionally substituted on

5 carbon by one or more R<sup>26</sup>;

R<sup>11</sup> and R<sup>12</sup> are independently selected from hydrogen, hydroxy, amino, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, *N*-(C<sub>1-4</sub>alkyl)amino, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl carbocyclylC<sub>1-4</sub>alkyl, heterocyclylC<sub>1-4</sub>alkyl; wherein R<sup>11</sup> and R<sup>12</sup> may be independently optionally substituted on carbon by one or more groups selected from R<sup>24</sup>; and wherein if said

10 heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>25</sup>;

R<sup>24</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, *N*-(C<sub>1-4</sub>alkyl)amino, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino,

15 C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl and C<sub>1-4</sub>alkylsulphonylamino;

R<sup>8</sup>, R<sup>10</sup>, R<sup>17</sup>, R<sup>19</sup> and R<sup>25</sup> are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, *N*-(C<sub>1-4</sub>alkyl)carbamoyl,

20 *N,N*-(C<sub>1-4</sub>alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl, heterocyclyl and phenylsulphonyl;

R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup> and R<sup>23</sup> are independently selected from hydrogen, phenyl, C<sub>1-4</sub>alkylsulphonyl and C<sub>1-4</sub>alkyl;

R<sup>26</sup> is selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, *N*-methyl-*N*-ethylamino, acetylamino, *N*-methylcarbamoyl, *N*-ethylcarbamoyl, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl,

25 *N*-methylsulphamoyl, *N*-ethylsulphamoyl, *N,N*-dimethylsulphamoyl, *N,N*-diethylsulphamoyl or *N*-methyl-*N*-ethylsulphamoyl;

or a pharmaceutically acceptable salt thereof;


in the manufacture of a medicament for use in the inhibition of 11βHSD1;

100804-2

- 7 -

with the proviso that said compound is not (1-methyl-1-pyrid-3-ylethyl)-(pyrid-3-yl)-ketone.

According to a further feature of the invention there is provided a compound of formula (Ia):



5

wherein:

**Ring A** is selected from furanyl, thienyl or pyridyl;

**R¹** is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; or two R¹ on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy group; wherein R¹ may be optionally substituted on carbon by one or more groups selected from R⁷; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R⁸;

15

n is 0-3; wherein the values of R¹ may be the same or different;

**R²** is selected from amino, C<sub>1-3</sub>alkoxy and N-(C<sub>1-3</sub>alkyl)amino; wherein R² may be optionally substituted on carbon by one or more groups selected from R⁹;

20 **Ring B** is 3-6 membered aryl or a 3-6 membered heteroaryl; wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R¹⁷;

**R⁶** is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein R⁶ may be optionally substituted on carbon by one or more groups selected from R¹⁸; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R¹⁹;

25

30

100804-2

- 8 -

$m$  is 0-3; wherein the values of  $R^6$  may be the same or different;


$Y$  is  $-S(O)_a-$ ,  $-O-$ ,  $-NR^{20}-$ ,  $-C(O)-$ ,  $-C(O)NR^{21}-$ ,  $-NR^{22}C(O)-$  or  $-SO_2NR^{23}-$ ; wherein  $a$  is 0 to 2;

$R^7$ ,  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N-(C_{1-4}alkyl)amino$ ,  $N,N-(C_{1-4}alkyl)_2amino$ ,  $C_{1-4}alkanoylamino$ ,  $N-(C_{1-4}alkyl)carbamoyl$ ,  $N,N-(C_{1-4}alkyl)_2carbamoyl$ ,  $C_{1-4}alkylS(O)_a$  wherein  $a$  is 0 to 2,  $C_{1-4}alkoxycarbonyl$ ,  $N-(C_{1-4}alkyl)sulphamoyl$ ,  $N,N-(C_{1-4}alkyl)_2sulphamoyl$ ,  $C_{1-4}alkylsulphonylamino$ , carbocyclyl and heterocyclyl;

$R^8$ ,  $R^{17}$  and  $R^{19}$  are independently selected from  $C_{1-4}alkyl$ ,  $C_{1-4}alkanoyl$ ,  $C_{1-4}alkylsulphonyl$ ,  $C_{1-4}alkoxycarbonyl$ , carbamoyl,  $N-(C_{1-4}alkyl)carbamoyl$ ,  $N,N-(C_{1-4}alkyl)_2carbamoyl$ , benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

$R^{20}$ ,  $R^{21}$ ,  $R^{22}$  and  $R^{23}$  are independently selected from hydrogen and  $C_{1-4}alkyl$ ; or a pharmaceutically acceptable salt thereof; with the proviso that said compound is not ( $\alpha$ -methoxybenzyl)-(pyrid-4-yl)-ketone, ( $\alpha$ -aminobenzyl)-(pyrid-3-yl)-ketone, [1-(fur-2-yl)-1-(ethoxy)methyl]-(fur-2-yl)-ketone or [1-(fur-2-yl)-1-(methoxy)methyl]-(fur-2-yl)-ketone.

According to a further feature of the invention there is provided a compound of formula (Ib):



wherein:

Ring A is thiazolyl;

$R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}alkyl$ ,  $C_{2-4}alkenyl$ ,  $C_{2-4}alkynyl$ ,  $C_{1-4}alkoxy$ ,  $C_{1-4}alkanoyl$ ,  $C_{1-4}alkanoyloxy$ ,  $N-(C_{1-4}alkyl)amino$ ,  $N,N-(C_{1-4}alkyl)_2amino$ ,  $C_{1-4}alkanoylamino$ ,  $N-(C_{1-4}alkyl)carbamoyl$ ,  $N,N-(C_{1-4}alkyl)_2carbamoyl$ ,  $C_{1-4}alkylS(O)_a$  wherein  $a$  is 0 to 2,  $C_{1-4}alkoxycarbonyl$ ,  $N-(C_{1-4}alkyl)sulphamoyl$ ,  $N,N-(C_{1-4}alkyl)_2sulphamoyl$ ,  $C_{1-4}alkylsulphonylamino$ , carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}alkylene-Y-$  and heterocyclyl $C_{0-4}alkylene-Y-$ ; or two  $R^1$  on

100804-2

- 9 -

adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy group; wherein R<sup>1</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>7</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>8</sup>;

n is 0-3; wherein the values of R<sup>1</sup> may be the same or different;

5 R<sup>2</sup> is selected from hydroxy, amino, C<sub>1-3</sub>alkoxy and N-(C<sub>1-3</sub>alkyl)amino; wherein R<sup>2</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>9</sup>;

Ring B is 3-6 membered aryl or a 3-6 membered heteroaryl; wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>17</sup>;

10 R<sup>6</sup> is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub>, wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, 15 N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein R<sup>6</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>18</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>19</sup>;

m is 0-3; wherein the values of R<sup>6</sup> may be the same or different;

20 Y is -S(O)<sub>x</sub>-, -O-, -NR<sup>20</sup>-, -C(O)-, -C(O)NR<sup>21</sup>-, -NR<sup>22</sup>C(O)- or -SO<sub>2</sub>NR<sup>23</sup>-; wherein a is 0 to 2;

R<sup>7</sup>, R<sup>9</sup> and R<sup>18</sup> are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, 25 N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub>, wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl;

R<sup>8</sup>, R<sup>17</sup> and R<sup>19</sup> are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, 30 C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;


R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup> and R<sup>23</sup> are independently selected from hydrogen and C<sub>1-4</sub>alkyl; or a pharmaceutically acceptable salt thereof.



100804-2

- 10 -

According to a further feature of the invention there is provided a compound of formula (Ic):



5 wherein:

Ring A is selected from furyl, thienyl, thiazolyl and pyridyl;

$R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}$ alkylene-Y- and heterocyclyl $C_{0-4}$ alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxy $C_{1-4}$ alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;

15  $n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

$R^2$  is selected from 3-6 membered aryl or carbon linked 3-6 membered heteroaryl; wherein  $R^2$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; and wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{10}$ ;

Ring B is 3-6 membered aryl or a carbon linked 3-6 membered heteroaryl; wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^6$  may be optionally substituted on carbon by one or more groups

100804-2

- 11 -

selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

$m$  is 0-3; wherein the values of  $R^6$  may be the same or different;

$Y$  is  $-S(O)_a-$ ,  $-O-$ ,  $-NR^{20}-$ ,  $-C(O)-$ ,  $-C(O)NR^{21}-$ ,  $-NR^{22}C(O)-$  or  $-SO_2NR^{23}-$ ; wherein  $a$  is

5 0 to 2;

$R^7$ ,  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N-(C_{1-4}alkyl)amino$ ,  $N,N-(C_{1-4}alkyl)_2amino$ ,  $C_{1-4}alkanoylamino$ ,  $N-(C_{1-4}alkyl)carbamoyl$ ,

10  $N,N-(C_{1-4}alkyl)_2carbamoyl$ ,  $C_{1-4}alkylS(O)_a$  wherein  $a$  is 0 to 2,  $C_{1-4}alkoxycarbonyl$ ,  $N-(C_{1-4}alkyl)sulphamoyl$ ,  $N,N-(C_{1-4}alkyl)_2sulphamoyl$ ,  $C_{1-4}alkylsulphonylamino$ , carbocyclyl and heterocyclyl;

$R^8$ ,  $R^{10}$ ,  $R^{17}$  and  $R^{19}$  are independently selected from  $C_{1-4}alkyl$ ,  $C_{1-4}alkanoyl$ ,  $C_{1-4}alkylsulphonyl$ ,  $C_{1-4}alkoxycarbonyl$ , carbamoyl,  $N-(C_{1-4}alkyl)carbamoyl$ ,

15  $N,N-(C_{1-4}alkyl)carbamoyl$ , benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;


$R^{20}$ ,  $R^{21}$ ,  $R^{22}$  and  $R^{23}$  are independently selected from hydrogen and  $C_{1-4}alkyl$ ;

or a pharmaceutically acceptable salt thereof;

with the proviso that said compound is not [1-(pyrazin-2-yl)-2-(2-fluorophenyl)ethyl]-(fur-2-yl)-ketone, [1-(pyrazin-2-yl)-2-(4-chlorophenyl)ethyl]-(fur-2-yl)-ketone, [2-(pyridin-3-yl)-1-

20 (2,4-dichlorophenyl)ethyl]-(pyrid-3-yl)-ketone, [2-(fur-2-yl)-1-(2,4-dichlorophenyl)ethyl]-(pyrid-3-yl)-ketone, [2-(4-nitrophenyl)-1-(2,4-dichlorophenyl)ethyl]-(pyrid-3-yl)-ketone, [2-(thien-2-yl)-1-(2,4-dichlorophenyl)ethyl]-(pyrid-3-yl)-ketone, [2-(phenyl)-1-(2,4-dichlorophenyl)ethyl]-(pyrid-3-yl)-ketone or [2-(4-chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyrid-3-yl)-ketone.

25 According to a further feature of the invention there is provided a compound of formula (Id):



wherein:

30 Ring A is thiazolyl;

100804-2

- 12 -

- $R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;
- 10  $n$  is 0-3; wherein the values of  $R^1$  may be the same or different;
- Ring B is 3-6 membered aryl or a 3-6 membered heteroaryl; wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;
- $R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;
- 20  $m$  is 0-3; wherein the values of  $R^6$  may be the same or different;
- Y is -S(O)<sub>a</sub>-, -O-, -NR<sup>20</sup>-, -C(O)-, -C(O)NR<sup>21</sup>-, -NR<sup>22</sup>C(O)- or -SO<sub>2</sub>NR<sup>23</sup>-; wherein a is 25 0 to 2;
- $R^7$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl;
- 30


100804-2

- 13 -

$R^8$ ,  $R^{17}$  and  $R^{19}$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

$R^{20}$ ,  $R^{21}$ ,  $R^{22}$  and  $R^{23}$  are independently selected from hydrogen and  $C_{1-4}$ alkyl;  
 5 or a pharmaceutically acceptable salt thereof;  
 with the proviso that said compound is not (phenethyl)-(5-aminothiazol-4-yl)-ketone.

According to a further feature of the invention there is provided a compound of formula (Ie):



(Ie)

10

wherein:

G is O or S;

$R^1$  is selected from fluoro, chloro, bromo, sulphamoyl, methyl, methoxy, ethoxy, acetyl or thiomethyl;

15  $n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

Ring B is 3-6 membered aryl or a 3-6 membered carbon linked heteroaryl; wherein if said heteroaryl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub>, wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

$m$  is 0-3; wherein the values of  $R^6$  may be the same or different;

$R^{18}$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,

30

100804-2

- 14 -

C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub>,  
wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl,

*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl;

**R<sup>17</sup>** and **R<sup>19</sup>** are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl,

5 C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, *N*-(C<sub>1-4</sub>alkyl)carbamoyl,

*N,N*-(C<sub>1-4</sub>alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

or a pharmaceutically acceptable salt thereof;


with the proviso that said compound is not (2,5-dimethylthien-3-yl)-(2,5-dimethylthien-3-

ylmethyl)-ketone; (2,5-dichlorothien-3-yl)-(benzyl)-ketone; (2,4,5-trichlorothien-3-yl)-

10 (benzyl)-ketone; (4-bromothien-3-yl)-(2-nitrobenzyl)-ketone; (2-methylfur-3-yl)-(benzyl)-

ketone; or (2,5-dimethylthien-3-yl)-(5-chlorothien-2-ylmethyl)-ketone.

According to a further feature of the invention there is provided a compound of  
formula (If):



(If)

15

wherein:

**R<sup>1</sup>** is selected from fluoro, chloro, bromo, sulphamoyl, methyl, methoxy, ethoxy,  
acetyl or thiomethyl;

**n** is 0-3; wherein the values of **R<sup>1</sup>** may be the same or different;

20 **R<sup>2</sup>** is *N*-(C<sub>1-4</sub>alkyl)amino; wherein **R<sup>2</sup>** may be optionally substituted on carbon by one  
or more groups selected from **R<sup>9</sup>**;

**R<sup>3</sup>** is selected from hydrogen or C<sub>1-4</sub>alkyl; wherein **R<sup>3</sup>** may be optionally substituted  
on carbon by one or more groups selected from **R<sup>9</sup>**;

25 **Ring B** is carbocyclyl or heterocyclyl; wherein if said heterocyclyl contains an -NH-  
moiety that nitrogen may be optionally substituted by a group selected from **R<sup>17</sup>**;

**R<sup>6</sup>** is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino,  
carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl,  
C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanyloxy, *N*-(C<sub>1-4</sub>alkyl)amino,  
*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl,  
30 *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub>, wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl,  
*N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl

100804-2

- 15 -

and heterocyclyl; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;


$m$  is 0-3; wherein the values of  $R^6$  may be the same or different;

- 5  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl, 10  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl and  $C_{1-4}$ alkylsulphonylamino;

$R^{17}$  and  $R^{19}$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt thereof;

- 15 with the proviso that said compound is not (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxyprop-2-ylamino)-4-methoxybenzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(butylamino)-4-methoxybenzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(ethylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxybut-2-ylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxybut-2-ylamino)-4-methoxybenzyl]-ketone; [3,4-dimethoxy-6-(methoxycarbonylmethyl)phenyl]-[ $\alpha$ -(methylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(butylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxyethylamino)-4-methoxybenzyl]-ketone; or (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxyethylamino)benzyl]-ketone.
- 20 (methylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(butylamino)benzyl]-ketone; (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxyethylamino)-4-methoxybenzyl]-ketone; or (4-methoxyphenyl)-[ $\alpha$ -(1-hydroxyethylamino)benzyl]-ketone.

According to a further feature of the invention there is provided a compound of formula (Ig):



(Ig)

wherein:

$R^1$  is selected from fluoro, chloro or methyl;

$R^2$  is  $C_{1-4}$ alkoxy; wherein  $R^2$  may be optionally substituted on carbon by one or more

- 30 groups selected from  $R^9$ ;

100804-2

- 16 -

$R^3$  is selected from hydrogen or  $C_{1-4}$ alkyl; wherein  $R^3$  may be optionally substituted on carbon by one or more groups selected from  $R^2$ ;

Ring B is carbocyclyl or a carbon linked heterocyclyl; wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  
5  $R^{17}$ ;

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  
10  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

15 m is 0-3; wherein the values of  $R^6$  may be the same or different;

$R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  
20  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl;


$R^{17}$  and  $R^{19}$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  
25  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt thereof;  
with the proviso that said compound is not (4-methylphenyl)-(α-methoxybenzyl)-ketone; (4-chlorophenyl)-(α-ethoxy-2-chlorobenzyl)-ketone; (4-chlorophenyl)-[1-(3-nitroimidazo[1,2-a]pyridin-8-yl)-1-(methoxy)methyl]-ketone; (4-methylphenyl)-(α-methoxy-α-methylbenzyl)-ketone; (2,4,6-trimethylphenyl)-(α-methoxy-α-methyl-2,4,6-trimethylbenzyl)-ketone; (2,4-dichlorophenyl)-(α-methoxybenzyl)-ketone; (4-fluorophenyl)-(α-methoxybenzyl)-ketone; (4-methylphenyl)-(α-methoxy-4-methylbenzyl)-ketone; (4-methylphenyl)-(α-*t*-butoxy-4-methylbenzyl)-ketone; (3-nitro-4-chlorophenyl)-(α-methoxy-3-nitro-4-chlorobenzyl)-ketone;  
30

100804-2

- 17 -

(4-methylphenyl)-( $\alpha$ -but-2-yloxybenzyl)-ketone; (4-chlorophenyl)-( $\alpha$ -isopropoxy-4-chlorobenzyl)-ketone; (4-chlorophenyl)-( $\alpha$ -isopropoxybenzyl)-ketone; (4-methylphenyl)-( $\alpha$ -isopropoxybenzyl)-ketone; (4-methylphenyl)-( $\alpha$ -isopropoxy-4-methylbenzyl)-ketone; (4-chlorophenyl)-( $\alpha$ -methoxybenzyl)-ketone; (4-chlorophenyl)-( $\alpha$ -methoxy-4-chlorobenzyl)-ketone; or (4-chlorophenyl)-( $\alpha$ -methoxy- $\alpha$ -methyl-4-chlorobenzyl)-ketone.

Accordingly to a further feature of the invention there is provided a compound of formula (Ib):



10 wherein:

Ring A is selected from furyl, thienyl, thiazolyl and pyridyl;

$R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ carbamoyl,  $C_{1-4}$ alkylS(O) $_a$  wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}$ alkylene-Y- and heterocyclyl $C_{0-4}$ alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxy $C_{1-4}$ alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;

$n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

$R^2$  and  $R^3$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino, carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}$ alkyl and heterocyclyl $C_{1-4}$ alkyl; wherein  $R^2$  and  $R^3$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{10}$ ;

$q$  is 0 or 1;

$p$  is 0 or 1;



100804-2

- 18 -

**Ring B** is a heterocyclyl linked to the sulphonyl of formula (II) via a nitrogen atom; wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclyl $C_{0-4}$ alkylene-Y- and heterocyclyl $C_{0-4}$ alkylene-Y-; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

m is 0-3; wherein the values of  $R^6$  may be the same or different;

$Y$  is -S(O)<sub>a</sub>-, -O-, -NR<sup>20</sup>-, -C(O)-, -C(O)NR<sup>21</sup>-, -NR<sup>22</sup>C(O)- or -SO<sub>2</sub>NR<sup>23</sup>-; wherein a is 0 to 2;

$R^7$ ,  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^7$ ,  $R^9$  and  $R^{18}$  may be independently optionally substituted on carbon by one or more  $R^{26}$ ;

$R^{24}$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl and  $C_{1-4}$ alkylsulphonylamino;

$R^8$ ,  $R^{10}$ ,  $R^{17}$ ,  $R^{19}$  and  $R^{25}$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

100804-2

- 19 -


$R^{20}$ ,  $R^{21}$ ,  $R^{22}$  and  $R^{23}$  are independently selected from hydrogen, phenyl and  $C_{1-4}$ alkyl;

$R^{26}$  is selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, *N*-methyl-*N*-ethylamino, acetylamino, *N*-methylcarbamoyl, *N*-ethylcarbamoyl, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, *N*-methylsulphamoyl, *N*-ethylsulphamoyl, *N,N*-dimethylsulphamoyl, *N,N*-diethylsulphamoyl or *N*-methyl-*N*-ethylsulphamoyl;

10 or a pharmaceutically acceptable salt thereof;

with the proviso that said compound is not (2-nitrofur-5-yl)-(morpholinosulphonylmethyl)-ketone.

According to a further feature of the invention there is provided a compound of formula (II):



15

(II)

wherein:

Ring A is selected from furyl, thienyl, thiazolyl and pyridyl;

$R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy, *N*-( $C_{1-4}$ alkyl)amino, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino, *N*-( $C_{1-4}$ alkyl)carbamoyl, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl, *N*-( $C_{1-4}$ alkyl)sulphamoyl, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;

25

n is 0-3; wherein the values of  $R^1$  may be the same or different;

$R^2$  and  $R^3$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, *N*-( $C_{1-4}$ alkyl)amino, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl,

30

100804-2

- 20 -

carbocyclylC<sub>1-4</sub>alkyl and heterocyclylC<sub>1-4</sub>alkyl; wherein R<sup>2</sup> and R<sup>3</sup> may be independently optionally substituted on carbon by one or more groups selected from R<sup>9</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>10</sup>;

5 q is 0 or 1;

p is 0 or 1;

Ring B is carbocyclyl or heterocyclyl; wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>17</sup>;

R<sup>6</sup> is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; wherein R<sup>6</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>18</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>19</sup>;

m is 0-3; wherein the values of R<sup>6</sup> may be the same or different;

20 Y is -S(O)<sub>2</sub>-, -O-, -NR<sup>20</sup>-, -C(O)-, -C(O)NR<sup>21</sup>-, -NR<sup>22</sup>C(O)- or -SO<sub>2</sub>NR<sup>23</sup>-; wherein a is 0 to 2;

R<sup>7</sup>, R<sup>9</sup> and R<sup>18</sup> are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, N-(C<sub>1-4</sub>alkyl)sulphamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-4</sub>alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein R<sup>7</sup>, R<sup>9</sup> and R<sup>18</sup> may be independently optionally substituted on carbon by one or more R<sup>26</sup>;

30 R<sup>24</sup> is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkanoyloxy, N-(C<sub>1-4</sub>alkyl)amino, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, N-(C<sub>1-4</sub>alkyl)carbamoyl, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub>

100804-2

- 21 -

wherein *a* is 0 to 2, C<sub>1-4</sub>alkoxycarbonyl, *N*-(C<sub>1-4</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl and C<sub>1-4</sub>alkylsulphonylamino;

*R*<sup>8</sup>, *R*<sup>10</sup>, *R*<sup>17</sup>, *R*<sup>19</sup> and *R*<sup>25</sup> are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl;

*R*<sup>16</sup>, *R*<sup>20</sup>, *R*<sup>21</sup>, *R*<sup>22</sup> and *R*<sup>23</sup> are independently selected from hydrogen, phenyl and C<sub>1-4</sub>alkyl;

*R*<sup>26</sup> is selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino, *N*-methyl-*N*-ethylamino, acetylamino, *N*-methylcarbamoyl, *N*-ethylcarbamoyl, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl, methylthio, ethylthio, methylsulphonyl, ethylsulphonyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, *N*-methylsulphamoyl, *N*-ethylsulphamoyl, *N,N*-dimethylsulphamoyl, *N,N*-diethylsulphamoyl or *N*-methyl-*N*-ethylsulphamoyl; or a pharmaceutically acceptable salt thereof.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, "C<sub>1-4</sub>alkyl" includes propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals therefore "carbocyclylC<sub>1-4</sub>alkyl" includes 1-carbocyclylpropyl, 2-carbocyclylethyl and 3-carbocyclylbutyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

"Heteroaryl" is a totally unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Suitably "heteroaryl" refers to a totally unsaturated, monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 8 - 10 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Examples and suitable values of the

100804-2

- 22 -

term "heteroaryl" are thienyl, furyl, thiazolyl, pyrazolyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyranyl, indolyl, pyrimidyl, pyrazinyl, pyridazinyl, benzothienyl, pyridyl and quinolyl. Particularly "heteroaryl" refers to thienyl, furyl, thiazolyl, pyridyl, benzothienyl, imidazolyl or pyrazolyl.

5 "3-6 Membered heteroaryl" is a totally unsaturated, mono or bicyclic ring containing 3-6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked. Suitably "3-6 membered heteroaryl" refers to a totally unsaturated, monocyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be  
10 carbon or nitrogen linked. Examples and suitable values of the term "3-6 membered heteroaryl" are thienyl, furyl, thiazolyl, pyrazolyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyranyl, pyrimidyl, pyrazinyl, pyridazinyl and pyridyl. Particularly "heteroaryl" refers to thienyl, furyl, thiazolyl, pyridyl, benzothienyl, imidazolyl or pyrazolyl.

15 "Aryl" is a totally unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms. Suitably "aryl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "aryl" include phenyl or naphthyl. Particularly "aryl" is phenyl.

"3-6 Membered aryl" is a totally unsaturated, mono or bicyclic carbon ring that contains 3-6 atoms. Suitably "3-6 membered aryl" is a monocyclic ring containing 5 or 6  
20 atoms. Suitable values for "3-6 membered aryl" include phenyl.

A "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally  
25 oxidised to form the S-oxides. Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 5 or 6 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)- or a ring sulphur atom may be optionally oxidised to form S-oxide(s). Examples and suitable values of  
30 the term "heterocyclyl" are thienyl, piperidiny, morpholiny, furyl, thiazolyl, pyridyl, imidazolyl, 1,2,4-triazolyl, thiomorpholiny, coumariny, pyrimidiny, phthalidyl, pyrazolyl, pyrazinyl, pyridazinyl, benzothienyl, benzimidazolyl, tetrahydrofuryl, [1,2,4]triazolo[4,3-a]pyrimidiny, piperidiny, indolyl, 1,3-benzodioxolyl and pyrrolidiny.

100804-2

- 23 -

A "carbocyclyl" is a saturated, partially saturated or unsaturated, mono or bicyclic carbon ring that contains 3-12 atoms; wherein a -CH<sub>2</sub>- group can optionally be replaced by a -C(O)-. Preferably "carbocyclyl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "carbocyclyl" include cyclopropyl, cyclobutyl, 1-oxocyclopentyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, phenyl, naphthyl, tetralinyl, indanyl or 1-oxoindanyl. Particularly "carbocyclyl" is cyclohexyl, phenyl, naphthyl or 2-6-dioxocyclohexyl.

- An example of "C<sub>1-4</sub>alkanoyloxy" is acetoxy. Examples of "C<sub>1-4</sub>alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C<sub>1-4</sub>alkoxy" include methoxy, ethoxy and propoxy. Examples of "oxyc<sub>1-4</sub>alkoxy" include oxymethoxy, oxyethoxy and oxypropoxy. Examples of "C<sub>1-4</sub>alkanoylamino" include formamido, acetamido and propionylamino. Examples of and "C<sub>1-4</sub>alkylS(O)<sub>*n*</sub>" wherein *n* is 0 to 2" include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl and ethylsulphonyl. Examples of and "C<sub>1-4</sub>alkylsulphonyl" include mesyl and ethylsulphonyl.
- Examples of "C<sub>1-4</sub>alkanoyl" include C<sub>1-3</sub>alkanoyl, propionyl and acetyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)amino" include methylamino and ethylamino. Examples of "*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino" include di-*N*-methylamino, di-(*N*-ethyl)amino and *N*-ethyl-*N*-methylamino. Examples of "C<sub>2-4</sub>alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C<sub>2-4</sub>alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)sulphamoyl" are *N*-(C<sub>1-3</sub>alkyl)sulphamoyl, *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl" are *N,N*-(dimethyl)sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "*N*-(C<sub>1-4</sub>alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "C<sub>1-4</sub>alkylsulphonylamino" are mesylamino and ethylsulphonylamino. Examples of "C<sub>0-4</sub>alkylene" are a direct bond, methylene and ethylene.

A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an

100804-2

- 24 -

organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomers and geometric isomers that possess 11 $\beta$ HSD1 inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula (I) that possess 11 $\beta$ HSD1 inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess 11 $\beta$ HSD1 inhibitory activity.

Particular values of variable groups are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

Ring A is selected from aryl.

Ring A is heteroaryl.

Ring A is selected from phenyl, naphthyl, thienyl, furyl, thiazolyl, pyridyl, benzothienyl, imidazolyl or pyrazolyl.

Ring A is selected from phenyl, naphthyl, thienyl, furyl, thiazolyl, pyridyl or imidazolyl.

Ring A is selected from phenyl, naphth-2-yl, thien-2-yl, thien-3-yl, fur-2-yl, thiazol-2-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, benzothien-3-yl, imidazol-2-yl or pyrazol-1-yl.

Ring A is selected from phenyl, naphth-2-yl, thien-2-yl, fur-2-yl, thiazol-2-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl or imidazol-2-yl.

Ring A is selected from phenyl, thien-2-yl, thien-3-yl, fur-2-yl, thiazol-2-yl, pyrid-2-yl, benzothien-3-yl, imidazol-2-yl or pyrazol-1-yl.

Ring A is phenyl substituted at the position para to the ketone.

R<sup>1</sup> is selected from halo, cyano, hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, N,N-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkylS(O)<sub>2</sub>, wherein a is 0, carbocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; or two R<sup>1</sup> on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy

100804-2

- 25 -

group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ;

Y is  $-S(O)_a-$  or  $-O-$ ; wherein a is 0 to 2; and

$R^7$  is halo.

- 5  $R^1$  is selected from halo, cyano, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $N,N$ -( $C_{1-6}$ alkyl) $_2$ amino,  $C_{1-6}$ alkylsulphonylamino, carbocyclyl and heterocyclyl $C_{0-6}$ alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxy $C_{1-6}$ alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ;

- 10 Y is  $-S(O)_a-$ , or  $-O-$ ; wherein a is 0 to 2; and

$R^7$  is halo.

$R^1$  is selected from fluoro, chloro, bromo, cyano, hydroxy, methyl, *t*-butyl, trifluoromethyl, methoxy, ethoxy, butoxy, dimethylamino, methylthio, 4-chlorophenyl, benzyloxy, morpholinosulphonyl and tetrahydrofur-2-yloxy; or two  $R^1$  on adjacent carbons

- 15 may form oxymethyleneoxy.

$R^1$  is selected from fluoro, chloro, bromo, iodo, cyano, hydroxy, methyl, pentyl, trifluoromethyl, methoxy, dimethylamino, methylsulphonylamino, phenyl, morpholinosulphonyl and tetrahydropyran-2-yloxy; or two  $R^1$  on adjacent carbons may form oxymethyleneoxy.

- 20  $R^1$  is selected from fluoro, chloro, bromo, cyano, methyl, trifluoromethyl, methoxy and ethoxy.

n is 0-2; wherein the values of  $R^1$  may be the same or different.

n is 0-1.

n is 0.

- 25 n is 1.

r is 1.

r is 2.

- $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl) $_2$ amino, carbocyclyl, heterocyclyl, carbocyclyl( $C_{1-4}$ alkyl), heterocyclyl( $C_{1-4}$ alkyl); wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently
- 30 optionally substituted on carbon by one or more groups selected from  $R^8$ ;



100804-2

- 26 -

$R^9$  is selected from halo, nitro, cyano, trifluoromethyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkoxycarbonyl and carbocyclyl; wherein  $R^9$  may be optionally substituted on carbon by one or more  $R^{26}$ ; wherein

$R^{26}$  is hydroxy.

- 5  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino, carbocyclyl, carbocyclyl( $C_{1-4}$ alkyl and heterocyclyl( $C_{1-4}$ alkyl; wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; wherein

$R^9$  is selected from halo, cyano,  $C_{1-4}$ alkyl and  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino.

- 10  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano, methyl, ethyl, propyl, isopropyl, ethoxy, isobutoxy, cyanomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminomethyl,  $N,N$ -dimethylaminomethyl,  $N,N$ -diethylaminomethyl,  $N,N$ -dipropylaminomethyl,  $N,N$ -diisopropylaminomethyl, 2-hydroxyethylaminomethyl, methylamino, ethylamino, propylamino, isopropylamino, 2-hydroxyethylamino, 2-( $N,N$ -diethylamino)ethylamino, 3-( $N,N$ -dimethylamino)propylamino,  $N,N$ -dipropylamino, phenyl, 2-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, piperidin-1-yl, indol-1-yl, 1,3-benzodioxol-5-yl, benzyl,  $\alpha$ -cyanobenzyl, 2-fluorobenzyl, 2-nitrobenzyl, 2-ethoxycarbonylbenzyl, 3-nitrobenzyl, 3-trifluoromethylbenzyl, 3-methoxycarbonylbenzyl, 4-fluorobenzyl, 4-chlorobenzyl, 4-nitrobenzyl, 4-methoxycarbonylbenzyl, 2,4-dichlorobenzyl, 3-nitro-6-methoxybenzyl, benzylamino, phenethylaminopyrrolidin-1-ylmethyl, piperidin-1-ylmethyl, morpholinomethyl, 5-nitrofur-2-ylmethyl, 2-methylthiazol-4-ylmethyl, 2-chlorothiazol-5-ylmethyl, pyrid-2-ylmethyl, pyrid-3-ylmethyl, pyrid-4-ylmethyl.

- 25  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, methyl, ethyl, cyanomethyl, diisopropylaminomethyl, methoxy, ethoxy, isopropoxy, ethylamino, isopropylamino, methylamino, phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl, benzyl, 4-chlorobenzyl, 2-fluorobenzyl, 4-fluorobenzyl and 2-chlorothiazol-5-ylmethyl.

- 30  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano, methyl, ethyl, propyl, isopropyl, ethoxy, cyanomethyl, methylamino, ethylamino, propylamino, isopropylamino, piperidin-1-yl, benzyl, 4-fluorobenzyl, 4-chlorobenzyl, 4-methoxycarbonylbenzyl, 2,4-dichlorobenzyl, benzylamino, piperidin-1-ylmethyl,

100804-2

- 27 -

morpholinomethyl, 2-methylthiazol-4-ylmethyl, 2-chlorothiazol-5-ylmethyl, pyrid-2-ylmethyl, pyrid-3-ylmethyl and pyrid-4-ylmethyl.

$R^2$  and  $R^3$  are not both methyl.

One of  $R^2$  and  $R^3$  is hydrogen.

- 5 One of  $R^2$  and  $R^3$  is selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}$ alkyl and heterocyclyl $C_{1-4}$ alkyl; and the other is selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}$ alkyl and heterocyclyl $C_{1-4}$ alkyl; wherein  $R^2$  and  $R^3$  may be
- 10 independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{10}$ .

$X$  is  $-S(O)_a-$ ,  $-O-$ ,  $-NR^{13}-$ ,  $-NR^{15}C(O)-$ ,  $-SO_2NR^{16}-$  or  $-NR^{16}SO_2-$ ; wherein  $a$  is 0 or 2;  $R^{13}$ ,  $R^{15}$  and  $R^{16}$  are independently selected from hydrogen, phenyl and  $C_{1-4}$ alkyl.

- 15  $X$  is  $-S(O)_a-$ ,  $-O-$ ,  $-NR^{13}-$ ,  $-NR^{15}C(O)-$ ,  $-SO_2NR^{16}-$  or  $-NR^{16}SO_2-$ ; wherein  $a$  is 0 or 2; and

$R^{13}$ ,  $R^{15}$  and  $R^{16}$  are independently selected from hydrogen, phenyl,  $C_{1-4}$ alkylsulphonyl and  $C_{1-4}$ alkyl.

$X$  is  $-S-$ ,  $-S(O)_2-$ ,  $-O-$ ,  $-NH-$ ,  $-NMe-$ ,  $-NHC(O)-$ ,  $-SO_2NMe-$  or  $-NPhSO_2-$ .

- 20  $X$  is  $-S-$ ,  $-S(O)_2-$ ,  $-O-$ ,  $-NMe-$ ,  $-NEt-$ ,  $-N(iPr)-$ ,  $-N(SO_2Me)-$ ,  $-NHC(O)-$ ,  $-NPhC(O)-$ ,  $-SO_2NH-$ ,  $-SO_2NMe-$ ,  $-SO_2NEt-$ ,  $-SO_2N(iPr)-$ ,  $-NMeSO_2-$ , or  $-NEtSO_2-$ .

$X$  is  $-S(O)_2-$ ,  $-O-$ ,  $-NH-$ ,  $-NMe-$ ,  $-NHC(O)-$ ,  $-SO_2NMe-$  or  $-NPhSO_2-$ .

$X$  is  $-SO_2NR^{16}-$ .

$X$  is  $-S(O)_a-$ ; wherein  $a$  is 2 and Ring B is a nitrogen linked heterocyclyl.

- 25  $q$  is 0.

$q$  is 1.

$p$  is 0.

$p$  is 1.

Ring B is carbocyclyl.

- 30 Ring B is heterocyclyl.

Ring B is phenyl, thien-2-yl, thien-3-yl, piperidin-1-yl, morpholino, morpholin-2-yl, 4-benzylmorpholin-2-yl, naphth-1-yl, naphth-2-yl, 2,6-dioxocyclohex-1-yl, cyclohexyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, imidazol-1-yl, 1-methylimidazol-2-yl, 1,2,4-triazol-1-

100804-2

- 28 -

yl, thiomorpholino, coumarin-7-yl, pyrimidin-2-yl, phthalid-3-yl, pyrazin-2-yl, pyridazin-3-yl, benzimidazol-1-yl or [1,2,4]triazolo[4,3-a]pyrimidin-5-yl.

$R^{17}$  is selected from  $C_{1-4}$ alkyl or benzyl.

Ring B is phenyl, thienyl, furyl, thiazolyl, piperidinyl, piperazinyl, morpholinyl, naphthyl, cyclohexyl, pyridyl, imidazolyl, 1,2,4-triazolyl, 1,3-benzodioxolyl, thiomorpholinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzimidazolyl or pyrimidinyl; wherein if Ring B contains an -NH- moiety, that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

$R^{17}$  is  $C_{1-4}$ alkyl or benzyl.

10 Ring B is phenyl, thienyl, piperidinyl, morpholinyl, naphthyl, 2,6-dioxocyclohexyl, cyclohexyl, pyridyl, imidazolyl, 1,2,4-triazolyl, thiomorpholinyl, coumarinyl, pyrimidinyl, phthalidyl, pyrazinyl, pyridazinyl, benzimidazolyl or [1,2,4]triazolo[4,3-a]pyrimidinyl; wherein if said imidazolyl or morpholinyl is linked via a carbon it may be optionally substituted on the -NH- by a group selected from  $R^{17}$ ;

15 Ring B is phenyl, thien-2-yl, fur-2-yl, thiazol-4-yl, thiazol-5-yl, thien-3-yl, piperidin-1-yl, 4-methylpiperazin-1-yl, morpholino, *N*-benzylmorpholin-1-yl, naphth-2-yl, cyclohexyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, imidazol-1-yl, 1-methylimidazol-2-yl, 1,2,4-triazol-1-yl, 1,3-benzodioxol-5-yl, thiomorpholino, pyrimidin-2-yl, pyrazin-2-yl, pyridazin-3-yl, benzimidazol-1-yl, benzimidazol-2-yl, 1-methylbenzimidazol-2-yl or pyrimidin-2-yl.

20 Ring B is phenyl, thien-2-yl, thien-3-yl, piperidin-1-yl, morpholino, morpholin-2-yl, 4-benzylmorpholin-2-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, thiomorpholino, pyrimidin-2-yl, phthalid-3-yl, pyrazin-2-yl, pyridazin-3-yl, benzimidazol-1-yl or [1,2,4]triazolo[4,3-a]pyrimidin-5-yl.

Ring B is phenyl substituted at the position para to  $-(CR^4R^5)_q$ .

25  $R^6$  is a substituent on carbon and is selected from halo, cyano, hydroxy, amino, carbamoyl, trifluoromethyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl, *N*-( $C_{1-4}$ alkyl)amino,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 or 2, carbocyclyl, heterocyclyl and heterocyclyl $C_{0-4}$ alkylene-Y-; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ;

Y is -S(O)<sub>2</sub>-;

30  $R^{18}$  is selected from halo, cyano, hydroxy, carbocyclyl and heterocyclyl.

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, carbamoyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino, *N*-( $C_{1-4}$ alkyl)carbamoyl, *N,N*-( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 or 2,

100804-2

- 29 -

C<sub>1-4</sub>alkoxycarbonyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, carbocyclyl, heterocyclyl and carbocyclylC<sub>0-4</sub>alkylene-Y-; wherein R<sup>6</sup> may be optionally substituted on carbon by one or more groups selected from R<sup>18</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>19</sup>;

5 Y is -C(O) or -C(O)NR<sup>21</sup>-;

R<sup>18</sup> is selected from halo, cyano, hydroxy, C<sub>1-4</sub>alkoxy and heterocyclyl;

R<sup>19</sup> is heterocyclyl; and

R<sup>21</sup> is hydrogen.

R<sup>6</sup> is a substituent on carbon and is selected from fluoro, chloro, bromo, cyano,

10 hydroxy, amino, carbamoyl, trifluoromethyl, methyl, *t*-butyl, cyanomethyl, methoxy, ethoxy, acetyl, 2-hydroxyethylamino, methylthio, mesyl, phenyl, 4-fluorophenyl, 2-thiazolin-2-yl, morpholinomethyl, and piperidin-1-ylsulphonyl.

R<sup>6</sup> is a substituent on carbon and is selected from fluoro, chloro, bromo, iodo, nitro, cyano, carbamoyl, methyl, propyl, isopropyl, butyl, *t*-butyl, hydroxymethyl, cyanomethyl,

15 morpholinomethyl, methoxy, ethoxy, 2-methoxyethoxy, acetyl, diethylamino, acetylamino, *N*-(isopropyl)carbamoyl, *N*-(isobutyl)carbamoyl, *N,N*-dimethylcarbamoyl, methoxymethylthio, methylthio, mesyl, methoxycarbonyl, ethoxycarbonyl, *N,N*-dimethylsulphamoyl, phenyl, cyclopentyl, 4-fluorophenyl, anilinocarbonyl, 4-(pyrid-4-yl)piperazin-1-yl, 2-thiazolin-2-yl, morpholino and 4-chlorobenzoyl.

20 R<sup>6</sup> is a substituent on carbon and is selected from fluoro, chloro, cyano, carbamoyl, trifluoromethyl, methyl, cyanomethyl, methoxy, ethoxy, acetyl, 2-hydroxyethylamino, mesyl, 4-fluorophenyl, 2-thiazolin-2-yl, morpholinomethyl and piperidin-1-ylsulphonyl.

m is 0-2; wherein the values of R<sup>6</sup> may be the same or different.

m is 0 or 1.

25 m is 0.

m is 1.

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

Ring A is selected from phenyl, naphthyl, thienyl, furyl, thiazolyl, pyridyl,

30 benzothienyl, imidazolyl or pyrazolyl;

R<sup>1</sup> is selected from halo, cyano, hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy,

*N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkylS(O)<sub>n</sub> wherein a is 0, carbocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; or two R<sup>1</sup> on adjacent carbons may form an oxyC<sub>1-4</sub>alkoxy

100804-2

- 30 -

group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ;

$Y$  is  $-S(O)_a-$  or  $-O-$ ; wherein  $a$  is 0 to 2; and

$R^7$  is halo;

5  $n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

$r$  is 1;

$R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N-(C_{1-4}$ alkyl)amino,  $N,N-(C_{1-4}$ alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}$ alkyl, heterocyclyl $C_{1-4}$ alkyl; wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently

10 optionally substituted on carbon by one or more groups selected from  $R^9$ ;

$R^9$  is selected from halo, nitro, cyano, trifluoromethyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N-(C_{1-4}$ alkyl)amino,  $N,N-(C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkoxycarbonyl and carbocyclyl; wherein  $R^9$  may be optionally substituted on carbon by one or more  $R^{26}$ ; wherein

$R^{26}$  is hydroxy;

15  $X$  is  $-S(O)_a-$ ,  $-O-$ ,  $-NR^{13}-$ ,  $-NR^{15}C(O)-$ ,  $-SO_2NR^{16}-$  or  $-NR^{16}SO_2-$ ; wherein  $a$  is 0 or 2;

$R^{13}$ ,  $R^{15}$  and  $R^{16}$  are independently selected from hydrogen, phenyl and  $C_{1-4}$ alkyl;

$q$  is 0 or 1;

$p$  is 0 or 1;

20 Ring B is phenyl, thienyl, piperidinyl, morpholinyl, naphthyl, 2,6-dioxocyclohexyl, cyclohexyl, pyridyl, imidazolyl, 1,2,4-triazolyl, thiomorpholinyl, coumarinyl, pyrimidinyl, phthalidyl, pyrazinyl, pyridazinyl, benzimidazolyl or [1,2,4]triazolo[4,3-*a*]pyrimidinyl; wherein if said imidazolyl or morpholinyl is linked via a carbon it may be optionally substituted on the  $-NH-$  by a group selected from  $R^{17}$ ;

$R^{17}$  is selected from  $C_{1-4}$ alkyl or benzyl;

25  $R^6$  is a substituent on carbon and is selected from halo, cyano, hydroxy, amino, carbamoyl, trifluoromethyl,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $N-(C_{1-4}$ alkyl)amino,  $C_{1-4}$ alkyl $S(O)_a$  wherein  $a$  is 0 or 2, carbocyclyl, heterocyclyl and heterocyclyl $C_{0-4}$ alkylene- $Y-$ ; wherein  $R^6$  may be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ;

$Y$  is  $-S(O)_2-$ ;

30  $R^{18}$  is selected from halo, cyano, hydroxy, carbocyclyl and heterocyclyl; and

$m$  is 0-3; wherein the values of  $R^6$  may be the same or different.

or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

100804-2

- 31 -

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

Ring A is selected from phenyl, naphth-2-yl, thien-2-yl, thien-3-yl, fur-2-yl, thiazol-2-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, benzothien-3-yl, imidazol-2-yl or pyrazol-1-yl;

5         $R^1$  is selected from fluoro, chloro, bromo, cyano, hydroxy, methyl, *t*-butyl, trifluoromethyl, methoxy, ethoxy, butoxy, dimethylamino, methylthio, 4-chlorophenyl, benzyloxy, morpholinosulphonyl and tetrahydrofur-2-yloxy; or two  $R^1$  on adjacent carbons may form oxymethyleneoxy;

$n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

10         $r$  is 1;

$R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano, methyl, ethyl, propyl, isopropyl, ethoxy, isobutoxy, cyanomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminomethyl, *N,N*-dimethylaminomethyl, *N,N*-diethylaminomethyl, *N,N*-dipropylaminomethyl, *N,N*-diisopropylaminomethyl, 2-  
15 hydroxyethylaminomethyl, methylamino, ethylamino, propylamino, isopropylamino, 2-hydroxyethylamino, 2-(*N,N*-diethylamino)ethylamino, 3-(*N,N*-dimethylamino)propylamino, *N,N*-dipropylamino, phenyl, 2-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, piperidin-1-yl, indol-1-yl, 1,3-benzodioxol-5-yl, benzyl,  $\alpha$ -cyanobenzyl, 2-fluorobenzyl, 2-nitrobenzyl, 2-  
20 ethoxycarbonylbenzyl, 3-nitrobenzyl, 3-trifluoromethylbenzyl, 3-methoxycarbonylbenzyl, 4-fluorobenzyl, 4-chlorobenzyl, 4-nitrobenzyl, 4-methoxycarbonylbenzyl, 2,4-dichlorobenzyl, 3-nitro-6-methoxybenzyl, benzylamino, phenethylaminopyrrolidin-1-ylmethyl, piperidin-1-ylmethyl, morpholinomethyl, 5-nitrofur-2-ylmethyl, 2-methylthiazol-4-ylmethyl, 2-chlorothiazol-5-ylmethyl, pyrid-2-ylmethyl, pyrid-3-ylmethyl, pyrid-4-ylmethyl;

25         $X$  is -S-, -S(O)<sub>2</sub>-, -O-, -NH-, -NMe-, -NHC(O)-, -SO<sub>2</sub>NMe- or -NPhSO<sub>2</sub>-;

$q$  is 0 or 1;

$p$  is 0 or 1;

Ring B is phenyl, thien-2-yl, thien-3-yl, piperidin-1-yl, morpholino, morpholin-2-yl, 4-benzylmorpholin-2-yl, naphth-1-yl, naphth-2-yl, 2,6-dioxocyclohex-1-yl, cyclohexyl,  
30 pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, imidazol-1-yl, 1-methylimidazol-2-yl, 1,2,4-triazol-1-yl, thiomorpholino, coumarin-7-yl, pyrimidin-2-yl, phthalid-3-yl, pyrazin-2-yl, pyridazin-3-yl, benzimidazol-1-yl or [1,2,4]triazolo[4,3-*a*]pyrimidin-5-yl;

100804-2

- 32 -

$R^6$  is a substituent on carbon and is selected from fluoro, chloro, bromo, cyano, hydroxy, amino, carbamoyl, trifluoromethyl, methyl, *i*-butyl, cyanomethyl, methoxy, ethoxy, acetyl, 2-hydroxyethylamino, methylthio, mesyl, phenyl, 4-fluorophenyl, 2-thiazolin-2-yl, morpholinomethyl, and piperidin-1-ylsulphonyl;

- 5  $m$  is 0-3; wherein the values of  $R^6$  may be the same or different;  
or a pharmaceutically acceptable salt thereof;  
in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

Therefore in a further aspect of the invention there is provided the use of a compound of formula (I) (as depicted above) wherein:

- 10 Ring A is selected from phenyl, naphthyl, thienyl, furyl, thiazolyl, pyridyl or imidazolyl.

- $R^1$  is selected from halo, cyano, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $N,N$ -( $C_{1-6}$ alkyl)<sub>2</sub>amino,  $C_{1-6}$ alkylsulphonylamino, carbocyclyl and heterocyclyl( $C_{0-6}$ alkylene-Y-; or two  $R^1$  on adjacent carbons may form an oxy $C_{1-4}$ alkoxy group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ;

Y is -S(O)<sub>a</sub>-, or -O-; wherein a is 0 to 2; and

$R^7$  is halo.

$n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

- 20  $r$  is 1 or 2;

$R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino, carbocyclyl, carbocyclyl $C_{1-4}$ alkyl and heterocyclyl $C_{1-4}$ alkyl; wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; wherein

- 25  $R^9$  is selected from halo, cyano,  $C_{1-4}$ alkyl and  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino.

X is -S(O)<sub>a</sub>-, -O-, -NR<sup>13</sup>-, -NR<sup>13</sup>C(O)-, -SO<sub>2</sub>NR<sup>16</sup>- or -NR<sup>16</sup>SO<sub>2</sub>-; wherein a is 0 or 2;

and

$R^{13}$ ,  $R^{15}$  and  $R^{16}$  are independently selected from hydrogen, phenyl,  $C_{1-4}$ alkylsulphonyl and  $C_{1-4}$ alkyl.

- 30  $q$  is 0 or 1.

$p$  is 0 or 1.

Ring B is phenyl, thienyl, furyl, thiazolyl, piperidinyl, piperazinyl, morpholinyl, naphthyl, cyclohexyl, pyridyl, imidazolyl, 1,2,4-triazolyl, 1,3-benzodioxolyl, thiomorpholinyl,

100804-2

- 33 -

pyrimidinyl, pyrazinyl, pyridazinyl, benzimidazolyl or pyrimidinyl; wherein if Ring B contains an -NH- moiety, that nitrogen may be optionally substituted by a group selected from R<sup>17</sup>;

R<sup>17</sup> is C<sub>1-4</sub>alkyl or benzyl.

- 5 R<sup>6</sup> is a substituent on carbon and is selected from halo, nitro, cyano, carbamoyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkanoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>amino, C<sub>1-4</sub>alkanoylamino, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-4</sub>alkylS(O)<sub>a</sub> wherein a is 0 or 2, C<sub>1-4</sub>alkoxycarbonyl, *N,N*-(C<sub>1-4</sub>alkyl)<sub>2</sub>sulphamoyl, carbocyclyl, heterocyclyl and carbocyclylC<sub>0-4</sub>alkylene-Y-; wherein R<sup>6</sup> may be optionally substituted on carbon by one or
- 10 more groups selected from R<sup>18</sup>; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R<sup>19</sup>;

Y is -C(O) or -C(O)NR<sup>21</sup>-;

R<sup>18</sup> is selected from halo, cyano, hydroxy, C<sub>1-4</sub>alkoxy and heterocyclyl;

R<sup>19</sup> is heterocyclyl; and

- 15 R<sup>21</sup> is hydrogen.

m is 0-3; wherein the values of R<sup>6</sup> may be the same or different;

or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for use in the inhibition of 11βHSD1.

- In another aspect of the present invention, a suitable compound of the invention, or a
- 20 pharmaceutically acceptable salt thereof, is selected from Group A:

(benzyl)-[4-(morpholinosulphonyl)phenyl]-ketone;

(2-methylpyrid-5-yloxymethyl)-(phenyl)-ketone;

[2-(3-chlorophenyl)-2-(1,2,4-triazol-1-yl)ethyl]-(phenyl)-ketone;

(4-chlorobenzyl)-(2-bromophenyl)-ketone;

- 25 (4-chlorobenzyl)-(3-bromophenyl)-ketone;

(3,4-dichlorobenzyl)-(3,4-dichlorophenyl)-ketone;

[α-(4-fluorobenzyl)benzyl]-(pyrid-3-yl)-ketone;

{α-[3-(*N,N*-dimethylamino)propylamino]benzyl}-(pyrid-3-yl)-ketone;

(2,4-dibromophenoxyethyl)-(phenyl)-ketone;

- 30 [α-(cyclohexylamino)-4-chlorobenzyl]-(4-chlorophenyl)-ketone;

[1-(cyclohexylamino)-1-(1,3-benzodioxol-5-yl)methyl]-(1,3-benzodioxol-5-yl)-ketone;

[α-(cyclohexylamino)-3,4-dimethoxybenzyl]-(2-chlorophenyl)-ketone;

[α-(cyclohexylamino)-4-methylbenzyl]-(4-methylphenyl)-ketone;



100804-2

- 34 -

- [ $\alpha$ -(cyclohexylamino)-2-chlorobenzyl]-(2-chlorophenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(*N,N*-dipropylaminomethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(*N,N*-diisopropylaminomethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(*N,N*-diethylaminomethyl)-4-methoxybenzyl]-(4-methoxyphenyl)-ketone;  
5 [  $\alpha$ -hydroxy- $\alpha$ -(*N,N*-diethylaminomethyl)-4-methylbenzyl]-(4-methylphenyl)-ketone;  
{ $\alpha$ -hydroxy- $\alpha$ -[2-(hydroxyethyl)aminomethyl]benzyl}-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(propylaminomethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(isopropylaminomethyl)benzyl]-(phenyl)-ketone;  
(phthalid-3-ylmethyl)-(4-chlorophenyl)-ketone;  
10 [2-(3-trifluoromethylphenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(4-fluorophenyl)-ketone;  
[2-(4-nitrophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(4-chlorophenyl)-ketone;  
[2-(2-fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(4-chlorophenyl)-ketone;  
[2-(2,4-dichlorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(phenyl)-ketone;  
(4-bromobenzyl)-(4-fluorophenyl)-ketone;  
15 [2-(4-fluorophenyl)-1-(pyrazin-2-yl)ethyl]-(phenyl)-ketone;  
(phthalid-3-ylmethyl)-(4-fluorophenyl)-ketone;  
[2-(2-fluorophenyl)-1-(pyrazin-2-yl)ethyl]-(fur-2-yl)-ketone;  
[2-(4-chlorophenyl)-1-(pyrid-3-yl)ethyl]-(4-chlorophenyl)-ketone;  
[2-(2,4-dichlorophenyl)-1-(pyridazin-3-yl)ethyl]-(phenyl)-ketone;  
20 [2-(4-chlorophenyl)-1-(pyridazin-3-yl)ethyl]-(phenyl)-ketone;  
[2-(4-chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyrid-3-yl)-ketone;  
[2-(4-chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(fur-2-yl)-ketone;  
(3,4-dichlorobenzyl)-(4-chlorophenyl)-ketone;  
(2-fluorobenzyl)-(4-chlorophenyl)-ketone;  
25 [2-(4-fluorophenyl)-1-(pyrazin-2-yl)ethyl]-(4-chlorophenyl)-ketone;  
[2-(1,2,4-triazol-1-yl)-3-methylbutyl]-(phenyl)-ketone;  
[2-(4-chlorophenyl)-1-(phenyl)ethyl]-(pyrid-3-yl)-ketone;  
[2-(2-fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(thien-2-yl)-ketone;  
[2-(phenyl)-1-(imidazol-1-yl)ethyl]-(4-chlorophenyl)-ketone;  
30 [1-methyl-1-(1,2,4-triazol-1-yl)ethyl]-(4-chlorophenyl)-ketone;  
[2-(2-aminophenylthio)-2-(4-methoxyphenyl)ethyl]-(4-methoxyphenyl)-ketone;  
[2-(2,4-dichlorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(2-chlorothien-5-yl)-ketone;  
[1-(hydroxy)-1-(thien-3-yl)methyl]-(thien-3-yl)-ketone;

100804-2

- 35 -

- ( $\alpha$ -hydroxybenzyl)-(4-*i*-butylphenyl)-ketone;  
[2-(4-chlorophenyl)-1-(4-methylphenyl)ethyl]-(pyrid-3-yl)-ketone;  
[(7-methyl[1,2,4]triazolo[4,3-*a*]pyrimidin-5-yl)oxymethyl]-(4-chlorophenyl)-ketone;  
(4-phenyl-2,6-dioxocyclohexylmethyl)-(4-bromophenyl)-ketone;  
5 ( $\alpha$ -ethoxy- $\alpha$ -ethylaminomethylbenzyl)-(phenyl)-ketone;  
[ $\alpha$ -(2-oxocyclopentyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(5-chloropyrimidin-2-yl)benzyl]-(phenyl)-ketone;  
(phenoxymethyl)-(3,5-dimethyl-2,3-dihydro-pyrazol-2-yl)-ketone;  
[2-(piperidin-1-yl)-1-(4-methylphenylsulphonyl)]-(phenyl)-ketone;  
10 (benzimidazol-1-ylmethyl)-(4-bromophenyl)-ketone;  
[[2-(2-hydroxyethylamino)-benzimidazol-1-yl]methyl]-(thien-2-yl)-ketone;  
(2-carbamoylphenoxymethyl)-(4-bromophenyl)-ketone;  
(morpholin-2-ylmethyl)-(phenyl)-ketone;  
(pyrimidin-2-ylsulphanylmethyl)-(4-bromophenyl)-ketone;  
15 (4-acetylbenzyl)-(4-chlorophenyl)-ketone;  
[(1-methylimidazol-2-yl)sulphanylmethyl]-(4-chlorophenyl)-ketone;  
(benzimidazol-1-ylmethyl)-(2,4-dichlorophenyl)-ketone;  
(4-methylbenzyl)-[4-(tetrahydropyran-2-yloxy)phenyl]-ketone;  
(phenylsulphonylmethyl)-(pyrid-2-yl)-ketone;  
20 (4-chlorophenoxymethyl)-(3,5-difluorophenyl)-ketone;  
[[1-(naphth-2-yl)-1-(hydroxy)methyl]-(4-dimethylaminophenyl)-ketone;  
( $\alpha$ -hydroxy-4-methoxybenzyl)-(naphth-2-yl)-ketone;  
(4-chlorophenethyl)-(2,4-difluorophenyl)-ketone;  
(4-fluorophenoxymethyl)-(4-chlorophenyl)-ketone;  
25 (phenoxymethyl)-(4-trifluoromethyl-2-fluorophenyl)-ketone;  
[1-methyl-1-(1,2,4-triazol-1-yl)ethyl]-(4-trifluoromethyl-2-fluorophenyl)-ketone;  
(4-fluorophenethyl)-(4-trifluoromethylphenyl)-ketone;  
(4-fluorophenethyl)-(2,4-difluorophenyl)-ketone;  
(4-fluorophenethyl)-(4-chlorophenyl)-ketone;  
30 (benzyl)-(3,4-dichlorophenyl)-ketone;  
[4-(piperidin-1-ylsulphonyl)phenoxymethyl]-(phenyl)-ketone;  
[2-(morpholinomethyl)-3,5-dimethylphenoxymethyl]-(phenyl)-ketone;  
(phenylsulphonylmethyl)-(3,4-dihydroxyphenyl)-ketone; and

- 36 -

(4-methylphenylsulphonylmethyl)-(4-chloro-3-methylphenyl)-ketone.

In a further aspect of the invention, there is provided the use of a compound or a pharmaceutically acceptable salt thereof, selected from Group B:

- (2,2-diphenylethyl)-(phenyl)-ketone;
- 5 (1,2-diphenylethyl)-(4-chlorophenyl)-ketone;
- (1-phenylpropyl)-(phenyl)-ketone;
- [2-(piperidin-1-yl)-1-(phenyl)ethyl]-(phenyl)-ketone;
- [2-(morpholino)-1-(phenyl)ethyl]-(phenyl)-ketone;
- [2-(dimethylamino)-1-(phenyl)ethyl]-(phenyl)-ketone;
- 10 [2-(phenyl)-1-(imidazol-1-yl)ethyl]-(phenyl)-ketone;
- (1,2-diphenylethyl)-(phenyl)-ketone;
- ( $\alpha$ -propylbenzyl)-(phenyl)-ketone;
- [ $\alpha$ -(cyanomethyl)benzyl]-(phenyl)-ketone;
- [N-(4-methylphenylsulphonyl)anilinomethyl]-(phenyl)-ketone;
- 15 (phenylsulphonylmethyl)-(phenyl)-ketone;
- [(1-methylimidazol-2-yl)sulphanylmethyl]-(4-bromophenyl)-ketone;
- (4-methylphenylsulphonylmethyl)-(4-bromophenyl)-ketone;
- (4-chlorophenylsulphonylmethyl)-(phenyl)-ketone;
- (4-chlorophenylsulphonylmethyl)-(phenyl)-ketone;
- 20 (phenylsulphonylmethyl)-(4-methoxyphenyl)-ketone;
- (phenylsulphonylmethyl)-(4-methylphenyl)-ketone;
- (4-methylphenylsulphonylmethyl)-(4-chlorophenyl)-ketone;
- (4-chlorophenylsulphonylmethyl)-(4-bromophenyl)-ketone;
- (benzylsulphonylmethyl)-(phenyl)-ketone;
- 25 (2-carbamoylphenoxyethyl)-(phenyl)-ketone;
- (naphth-2-yloxyethyl)-(phenyl)-ketone;
- (phenoxyethyl)-(phenyl)-ketone;
- (4-chlorophenoxyethyl)-(phenyl)-ketone;
- (phenoxyethyl)-(4-chlorophenyl)-ketone;
- 30 (4-cyanophenoxyethyl)-(phenyl)-ketone;
- (4-*t*-butylphenoxyethyl)-(4-chlorophenyl)-ketone;
- (N-methylanilinomethyl)-(phenyl)-ketone;
- (4-chlorobenzamidomethyl)-(4-bromophenyl)-ketone;

100804-2

- 37 -

- [1-(cyano)-1-(thien-2-yl)ethyl]-(phenyl)-ketone;  
(phenethyl)-(4-bromophenyl)-ketone;  
[2-(2-methoxyphenyl)ethyl]-(phenyl)-ketone;  
(2-(cyano)-2-(phenyl)ethyl)-(phenyl)-ketone;  
5 (phenethyl)-(phenyl)-ketone;  
(phenethyl)-(2-methoxyphenyl)-ketone;  
(3,4-dimethoxyphenethyl)-(phenyl)-ketone;  
(phenethyl)-(4-chlorophenyl)-ketone;  
( $\alpha$ -hydroxybenzyl)-(phenyl)-ketone;  
10 ( $\alpha$ -hydroxy-4-chlorobenzyl)-(4-chlorophenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(*N,N*-diethylaminomethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(piperidin-1-ylmethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(*N,N*-dimethylaminomethyl)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -hydroxy- $\alpha$ -(morpholinomethyl)benzyl]-(phenyl)-ketone;  
15 ( $\alpha$ -hydroxy-4-chlorobenzyl)-(4-methoxyphenyl)-ketone;  
( $\alpha$ -ethoxybenzyl)-(phenyl)-ketone;  
( $\alpha$ -hydroxy- $\alpha$ -ethylbenzyl)-(phenyl)-ketone;  
( $\alpha$ -hydroxybenzyl)-(4-methoxyphenyl)-ketone;  
[1-(1,2,4-triazol-1-yl)-1-(ethoxy)methyl]-(4-chlorophenyl)-ketone;  
20 [1-(thien-2-yl)-1-(hydroxy)methyl]-(thien-2-yl)-ketone;  
( $\alpha$ -hydroxybenzyl)-(4-methoxyphenyl)-ketone;  
( $\alpha$ -hydroxy-4-methoxybenzyl)-(phenyl)-ketone;  
( $\alpha$ -isopropoxybenzyl)-(phenyl)-ketone;  
( $\alpha$ -isobutoxybenzyl)-(phenyl)-ketone;  
25 ( $\alpha$ -aminobenzyl)-(4-chlorophenyl)-ketone;  
( $\alpha$ -[2-(*N,N*-diethylamino)ethylamino]benzyl)-(phenyl)-ketone;  
( $\alpha$ -isopropylaminobenzyl)-(phenyl)-ketone;  
[ $\alpha$ -(piperidin-1-yl)-4-chlorobenzyl]-(4-chlorophenyl)-ketone;  
[ $\alpha$ -(benzylamino)benzyl]-(phenyl)-ketone;  
30 [ $\alpha$ -(4-chloroanilino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(cyclohexylamino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(*N,N*-dipropylamino)benzyl]-(phenyl)-ketone;

100804-2

- 38 -

- [ $\alpha$ -(2-hydroxyethylamino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(phenethylamino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(ethylamino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(propylamino)benzyl]-(phenyl)-ketone;  
5 [  $\alpha$ -(methylamino)benzyl]-(phenyl)-ketone;  
[ $\alpha$ -(anilino)benzyl]-(fur-2-yl)-ketone;  
[1-(benzimidazol-1-yl)-1-(anilino)methyl]-(phenyl)-ketone;  
(4-chlorobenzyl)-(phenyl)-ketone;  
(benzyl)-(4-ethoxyphenyl)-ketone;  
10 (4-methoxybenzyl)-(4-methoxyphenyl)-ketone;  
(benzyl)-(4-methylphenyl)-ketone;  
[4-(benzyl)morpholin-2-ylmethyl]-(phenyl)-ketone;  
(pyrid-2-ylmethyl)-(4-chlorophenyl)-ketone;  
(2-chlorobenzyl)-(4-chlorophenyl)-ketone;  
15 (4-chlorobenzyl)-(4-chlorophenyl)-ketone;  
(pyrid-3-ylmethyl)-(4-chlorophenyl)-ketone;  
(4-bromobenzyl)-(4-chlorophenyl)-ketone;  
(2,4-dichlorobenzyl)-(4-chlorophenyl)-ketone;  
(4-chlorobenzyl)-(4-methylphenyl)-ketone;  
20 (4-chlorobenzyl)-(4-bromophenyl)-ketone;  
(benzyl)-(2-chlorophenyl)-ketone;  
(4-methoxybenzyl)-(phenyl)-ketone;  
( $\alpha$ -methylbenzyl)-(phenyl)-ketone;  
(benzyl)-[4-(4-chlorophenyl)phenyl]-ketone;  
25 (4-fluorobenzyl)-(4-bromophenyl)-ketone;  
(4-chlorobenzyl)-(4-methoxyphenyl)-ketone;  
(4-methylbenzyl)-(4-methoxyphenyl)-ketone;  
(pyrid-2-ylmethyl)-(phenyl)-ketone;  
( $\alpha,\alpha$ -dimethylbenzyl)-(phenyl)-ketone;  
30 (4-methylbenzyl)-(pyrid-3-yl)-ketone;  
(pyrid-4-ylmethyl)-(pyrid-4-yl)-ketone;  
(4-methoxybenzyl)-(4-bromophenyl)-ketone;  
(4-methylthiobenzyl)-(4-fluorophenyl)-ketone;

- 39 -

- (benzyl)-(4-benzyloxyphenyl)-ketone;  
(4-fluorobenzyl)-(4-fluorophenyl)-ketone;  
( $\alpha$ -methylbenzyl)-(phenyl)-ketone;  
(4-methoxybenzyl)-(4-fluorophenyl)-ketone;  
5 (thiomorpholinomethyl)-(thianaphthen-3-yl)-(phenyl)-ketone;  
(benzyl)-(4-butoxyphenyl)-ketone;  
(2,2-diphenylethyl)-(2,4,6-trimethylphenyl)-ketone;  
[2-(2-hydroxyphenyl)-2-phenylethyl]-(phenyl)-ketone;  
(cyclohexylmethyl)-(phenyl)-ketone;  
10 (benzyl)-(2-bromothien-5-yl)-ketone;  
(1,2-diphenyl-2-cyanoethyl)-(phenyl)-ketone;  
(4-methoxybenzyl)-(3-bromophenyl)-ketone;  
( $\alpha$ -hydroxybenzyl)-(3-methoxyphenyl)-ketone;  
[ $\alpha$ -(pyrrolidin-1-ylmethyl)benzyl]-(phenyl)-ketone;  
15 [ $\alpha$ -(pyridin-2-ylamino)-4-methoxybenzyl]-(4-methoxyphenyl)-ketone;  
(4-chlorobenzyl)-(4-fluorophenyl)-ketone;  
(benzyl)-[4-(tetrahydropyran-2-yloxy)phenyl]-ketone;  
(4-chlorophenylsulphonylmethyl)-(4-chlorophenyl)-ketone;  
(4-methyl- $\alpha$ -hydroxybenzyl)-(4-chlorophenyl)-ketone;  
20 (4-methylbenzyl)-(4-chlorophenyl)-ketone;  
(4-fluoro- $\alpha$ -hydroxybenzyl)-(4-fluorophenyl)-ketone;  
(4-methoxy- $\alpha$ -hydroxybenzyl)-(4-methoxyphenyl)-ketone;  
( $\alpha$ -methyl- $\alpha$ -hydroxybenzyl)-(phenyl)-ketone;  
(1-methyl-1-morpholinoethyl)-(4-methylsulphanylphenyl)-ketone;  
25 [2-(phenylsulphonyl)-2-(phenyl)ethyl]-(phenyl)-ketone  
(1,3-diphenylprop-2-yl)-(phenyl)-ketone;  
(naphth-1-yloxymethyl)-(phenyl)-ketone;  
(phenoxymethyl)-(4-methylphenyl)-ketone;  
(4-methylcoumarin-7-yloxymethyl)-(4-methoxyphenyl)-ketone;  
30 (imidazol-1-ylmethyl)-(2-chlorothien-5-yl)-ketone;  
(thien-2-ylsulphonylmethyl)-(4-chlorophenyl)-ketone;  
(1-methylimidazol-2-ylsulphanylmethyl)-(3,4-difluorophenyl)-ketone;  
(1-methylimidazol-2-ylsulphonylmethyl)-(4-chlorophenyl)-ketone;

100804-2

- 40 -

- (3-trifluoromethylpyrid-6-ylsulphonylmethyl)-(4-chlorophenyl)-ketone;  
(4-methyl- $\alpha$ -hydroxybenzyl)-(4-methylphenyl)-ketone;  
(4-bromophenoxyethyl)-(phenyl)-ketone;  
(4-ethoxyanilinomethyl)-(4-methylphenyl)-ketone; and  
5 (2,4,6-trichlorophenoxyethyl)-(phenyl)-ketone;  
in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

In another aspect of the present invention, a suitable compound of the invention, or a pharmaceutically acceptable salt thereof, is selected from Group C:

- (phenyl)-[2-(piperidin-1-yl)-2-(2-chlorophenyl)ethyl]-ketone;  
10 (phenyl)-(2,4,5-trichlorophenoxyethyl)-ketone;  
(phenyl)-[ $\alpha$ -hydroxy- $\alpha$ -(butylaminomethyl)benzyl]-ketone;  
(4-fluorophenyl)-[2-(3,4-dichlorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(thien-2-yl)-[2-(4-fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(phenyl)-[2-(3-nitrophenyl)-2-(2-aminophenylthio)ethyl]-ketone;  
15 (pyrid-3-yl)-[ $\alpha$ -(*t*-butanonylmethyl)-4-chlorobenzyl]-ketone;  
(4-methoxyphenyl)-[4-(4-fluorobenzoyl)piperidin-1-ylmethyl]-ketone;  
(phenyl)-(2-nitro-4-chlorophenoxyethyl)-ketone;  
(phenyl)-(2,6-dibromo-3-ethoxycarbonylphenoxyethyl)-ketone;  
(4-bromophenyl)-(4-nitrophenoxyethyl)-ketone; and  
20 (phenyl)-(4-nitrophenoxyethyl)-ketone.

In a further aspect of the invention, there is provided the use of a compound or a pharmaceutically acceptable salt thereof, selected from Group D:

- (phenyl)-(benzoyl)-ketone;  
(phenyl)-[2-(4-methoxyphenyl)-2-cyanoethyl]-ketone;  
25 (phenyl)-[2-(phenyl)-2-(2-methoxyethylthio)ethyl]-ketone;  
(4-methylphenyl)-(4-methylbenzoyl)-ketone;  
(phenyl)-[2-(2-chlorophenyl)-2-cyanoethyl]-ketone;  
(phenyl)-(4-methylphenylsulphonylmethyl)-ketone;  
(4-chlorophenyl)-[2-(phenyl)-2-cyanoethyl]-ketone;  
30 (phenyl)-[2-(pyrrolidin-1-yl)-2-(phenyl)ethyl]-ketone;  
(4-bromophenyl)-[2-(piperidin-1-yl)-2-(phenyl)ethyl]-ketone;  
(phenyl)-[ $\alpha$ -(allylamino)benzyl]-ketone;  
(phenyl)-[ $\alpha$ -phenyl- $\alpha$ -hydroxybenzyl]-ketone;

100804-2

- 41 -

- (phenyl)-[2-(3-methoxyanilino)-2-(phenyl)ethyl]-ketone;  
(phenyl)-[ $\alpha$ -phenyl- $\alpha$ -hydroxy-4-methoxybenzyl]-ketone;  
(4-fluorophenyl)-[2-(4-chlorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(4-chlorophenyl)-[2-(4-cyanophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;
- 5 (phenyl)-[2-(morpholino)-2-(phenyl)ethyl]-ketone;  
(4-fluorophenyl)-[2-(2-fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(phenyl)-[2-(4-fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(phenyl)-[2-(phenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(4-chlorophenyl)-[2-(phenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;
- 10 (phenyl)-(benzylsulphinylmethyl)-ketone;  
(5-chlorothiophen-2-yl)-[2-(3,4-dichlorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(phenyl)-[2-(cyano)-2-(4-chlorophenyl)ethyl]-ketone;  
(thien-2-yl)-[2-(phenyl)-1-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(4-hydroxyphenyl)-(benzoyl)-ketone;
- 15 (phenyl)-[2-(morpholino)-1-(benzyl)ethyl]-ketone;  
(phenyl)-[2-(2-methoxyphenyl)-2-(2-aminophenylthio)ethyl]-ketone;  
(phenyl)-[2-(1,3-benzodioxol-5-yl)-2-(2-aminophenylthio)ethyl]-ketone;  
(phenyl)-[2-(4-fluorobenzoyl)-2-(1,2,4-triazol-1-yl)ethyl]-ketone;  
(3,4-dimethylphenyl)-[4-(4-fluorobenzoyl)piperidin-1-ylmethyl]-ketone;
- 20 (4-methoxyphenyl)-( $\alpha$ -methylbenzyl)-ketone;  
(4-methoxyphenyl)-(3-methylbenzyl)-ketone;  
(3-methyl-4-methoxyphenyl)-(benzyl)-ketone;  
(4-fluorophenyl)-(benzimidazol-1-ylmethyl)-ketone;  
(phenyl)-(1-methyl-1-imidazol-1-ylethyl)-ketone;
- 25 (phenyl)-(2-methylaminobenzimidazol-1-ylmethyl)-ketone;  
(4-chlorophenyl)-(2,4-dichlorophenoxyethyl)-ketone;  
(4-chlorophenyl)-(2,4,6-trichlorophenoxyethyl)-ketone;  
(4-bromophenyl)-[2-(trifluoromethyl)benzoylaminomethyl]-ketone;  
(4-bromophenyl)-( $\alpha$ -homopiperidin-1-ylbenzyl)-ketone;
- 30 (4-chlorophenyl)-(4-chloroanilinomethyl)-ketone;  
(phenyl)-[N-(benzoyl)anilinomethyl]-ketone;  
(phenyl)-(3-methylindol-1-ylmethyl)-ketone;  
(phenyl)-(2,4-dichlorobenzoylaminomethyl)-ketone;



100804-2

- 42 -

(phenyl)-[2-(phenyl)-1-(ethoxycarbonyl)ethyl]-ketone;

(4-chlorophenyl)-(4-chlorobenzoylaminomethyl)-ketone; and

(4-chlorophenyl)-(2-fluorobenzoylaminomethyl)-ketone;


in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1.

- 5 In another aspect of the invention, preferred compounds of the invention are any one of the Examples or a pharmaceutically acceptable salt thereof.

In another aspect of the invention, preferred compounds of the invention are any one of the Reference Examples or a pharmaceutically acceptable salt thereof.

- Another aspect of the present invention provides a process for preparing a compound  
10 of formula (I) or a pharmaceutically acceptable salt thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of:

*Process 1*): reacting a compound of formula (II):



(II)

- 15 wherein V is a displaceable group; with an organometallic reagent of formula (III):




(III)

wherein M is a metal reagent;

*Process 2*): for compounds of formula (I) wherein r is 1 and one of R<sup>2</sup> and R<sup>3</sup> is hydroxy;

- 20 reacting a compound of formula (IV):



(IV)

with a compound of formula (V) or (VI):




25

wherein M is a metal reagent;

100804-2

- 43 -

*Process 3*): for compounds of formula (I) wherein one of  $R^4$  and  $R^5$  is hydroxy; reacting a compound of formula (VII):



(VII)

5 with a compound of formula (VIII) or (IX):




(VIII)



(IX)

wherein M is a metal reagent;


*Process 4*): for compounds of formula (I) wherein p is 0, q is 1, r is 1 and  $R^3$  and  $R^5$  are  
10 hydrogen; hydrogenating compound of formula (X):



(X)


(or its corresponding Z isomer);

*Process 5*) for compounds of formula (I) wherein p is 1, X is  $-SO_2-$ , r is 1 and q is 0; reacting  
15 a compound of formula (XI):



(XI)

with a compound of formula (XII):



(XII)

20

100804-2


- 44 -

*Process 6)* for compounds of formula (I) wherein Ring A is a nitrogen linked heteroaryl;  
 reacting a compound of formula (II), wherein V is hydroxy, or an activated derivative thereof  
 forming an activated acid, with a compound of formula (XIII):




(XIII)

*Process 7)* for compounds of formula (I) wherein X is  $-C(O)NR^{14}-$ ; reacting an acid of  
 formula (XIV):




(XIV)

10 or an activated derivative thereof; with an amine of formula (XV):




(XV)

*Process 8)* for compounds of formula (I) wherein X is  $-NR^{15}C(O)-$ ; reacting an amine of  
 formula (XVI):



(XVI)

with an acid of formula (XVII):




(XVII)

20 or an activated derivative thereof;


100804-2

- 45 -


*Process 9)* for compounds of formula (I) wherein X is  $-\text{SO}_2\text{NR}^{16}-$ ; reacting a compound of formula (XVIII):



5 wherein L is a displaceable group; with an amine of formula (XIX):




*Process 10)* for compounds of formula (I) wherein X is  $-\text{NR}^{16}\text{SO}_2-$ ; reacting an amine of formula (XX):




10

with a compound of formula (XXI):



15 wherein L is a displaceable group;

*Process 11)* for compounds of formula (I) wherein X is  $-\text{O}-$ ,  $-\text{NR}^{13}-$  or  $-\text{S}-$ ; reacting a compound of formula (XX):




20

100804-2

- 46 -


wherein V is -OH, -NR<sup>13</sup>H or -SH; with a compound of formula (XXI):



(XXI)


wherein L is a displaceable group;

- 5 *Process 12)* for compounds of formula (I) wherein X is -O-, -NR<sup>13</sup>- or -S-; reacting a compound of formula (XX):



(XX)

wherein L is a displaceable group; with a compound of formula (XXI):



(XXI)

wherein V is -OH, -NR<sup>13</sup>H or -SH;

and thereafter if necessary or desirable:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- 15 ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt thereof.

L is a displaceable group, suitable values for L include halo, particularly chloro or bromo, or mesyloxy.

V is a displaceable group, suitable values for V include the Weinreb amide *N*-methyl-  
20 *N*-methoxyamine.

M is a metal reagent. Suitable values for M include Grignard reagents such as MgBr and lithium.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and active esters, for example pentafluorophenyl esters.

100804-2

- 47 -

The reactions described above may be performed under standard conditions. The intermediates described above are commercially available, are known in the art or may be prepared by known procedures.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, *Protective Groups in Organic Synthesis*, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali

100804-2

- 48 -

metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for  
5 example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for  
10 example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide.  
15 Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed,  
20 for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

25 As stated hereinbefore the compounds defined in the present invention possess 11 $\beta$ HSD1 inhibitory activity. These properties may be assessed using the following assay.

#### Assay

HeLa cells (human cervical carcinoma derived cells) were stably transfected with a construct containing four copies of the glucocorticoid response element (GRE) linked to a  
30 beta-galactosidase reporter gene (3 kb lac Z gene derived from pSV-B-galactosidase). These cells were then further stably transfected with a construct containing full-length human 11 $\beta$ HSD1 enzyme (in pCMVHyg) to create GRE4- $\beta$ Gal/11 $\beta$ HSD1 cells. The principal of the assay is as follows. Cortisone is freely taken up by the cells and is converted to cortisol by

100804-2

- 49 -

11 $\beta$ HSD1 oxo-reductase activity and cortisol (but not cortisone) binds to and activates the glucocorticoid receptor. Activated glucocorticoid receptor then binds to the GRE and initiates transcription and translation of  $\beta$ -galactosidase. Enzyme activity can then be assayed with high sensitivity by colourimetric assay. Inhibitors of 11 $\beta$ HSD1 will reduce the conversion of  
5 cortisone to cortisol and hence decrease the production of  $\beta$ -galactosidase.

Cells were routinely cultured in DMEM (Invitrogen, Paisley, Renfrewshire, UK) containing 10% foetal calf serum (LabTech), 1% glutamine (Invitrogen), 1% penicillin & streptomycin (Invitrogen), 0.5 mg/ml G418 (Invitrogen) & 0.5mg/ml hygromycin (Boehringer). Assay media was phenol red free-DMEM containing 1% glutamine, 1%  
10 penicillin & streptomycin.

Compounds (1mM) to be tested were dissolved in dimethyl sulphoxide (DMSO) and serially diluted into assay media containing 10% DMSO. Diluted compounds were then plated into transparent flat-bottomed 384 well plates (Matrix, Hudson NH, USA).

The assay was carried out in 384 well microtitre plate (Matrix) in a total volume of  
15 50 $\mu$ l assay media consisting of cortisone (Sigma, Poole, Dorset, UK, 1 $\mu$ M), HeLa GRE4- $\beta$ Gal/11 $\beta$ HSD1 cells (10,000 cells) plus test compounds (3000 to 0.01 nM). The plates were then incubated in 5% O<sub>2</sub>, 95% CO<sub>2</sub> at 37°C overnight.

The following day plates were assayed by measurement of  $\beta$ -galactosidase production.

A cocktail (25 $\mu$ l) consisting of 10X Z-buffer (600 mM Na<sub>2</sub>HPO<sub>4</sub>, 400 mM  
20 NaH<sub>2</sub>PO<sub>4</sub>.2H<sub>2</sub>O, 100 mM KCl, 10 mM MgSO<sub>4</sub>.7H<sub>2</sub>O, 500 mM  $\beta$ -mercaptoethanol, pH 7.0), SDS (0.2%), chlorophenol red- $\beta$ -D-galactopyranoside (5mM, Roche Diagnostics) was added per well and plates incubated at 37°C for 3-4hours.  $\beta$ -Galactosidase activity was indicated by a yellow to red colour change (absorbance at 570nm) measured using a Tecan Spectrafluor Ultra.

25 The calculation of median inhibitory concentration (IC<sub>50</sub>) values for the inhibitors was performed using Origin 6.0 (Microcal Software, Northampton MA USA). Dose response curves for each inhibitor were plotted as OD units at each inhibitor concentration with relation to a maximum signal (cortisone, no compound) and IC<sub>50</sub> values calculated. Compounds of the present invention typically show an IC<sub>50</sub> <10 $\mu$ M.

30 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih) or (Ii) or a pharmaceutically acceptable salt thereof, or a compound selected from Group A, Group C or the Examples, or a pharmaceutically acceptable salt thereof, as defined



hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical  
5 administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

The compound of formula (I), or a pharmaceutically acceptable salt thereof, will normally be administered to a warm-blooded animal at a unit dose within the range 0.1 –  
10 50 mg/kg that normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-1000 mg of active ingredient. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

15 We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt thereof, are effective 11 $\beta$ HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndromes.

It is to be understood that where the term "metabolic syndrome" is used herein, this relates to metabolic syndrome as defined in 1) and/or 2) or any other recognised definition of  
20 this syndrome. Synonyms for "metabolic syndrome" used in the art include Reaven's Syndrome, Insulin Resistance Syndrome and Syndrome X. It is to be understood that where the term "metabolic syndrome" is used herein it also refers to Reaven's Syndrome, Insulin Resistance Syndrome and Syndrome X.

According to a further aspect of the present invention there is provided a compound of  
25 the formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih) or (Ii) or a pharmaceutically acceptable salt thereof, or a compound selected from Group A, Group C or the Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

Thus according to this aspect of the invention there is provided a compound of the  
30 formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih) or (Ii) or a pharmaceutically acceptable salt thereof, or a compound selected from Group A, Group C or the Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore for use as a medicament.

100804-2

- 51 -

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an  $11\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man.

- 5        According to another feature of the invention there is provided the use of a compound of the formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih) or (II) or a pharmaceutically acceptable salt thereof, or a compound selected from Group A, Group C or the Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an  $11\beta$ HSD1 inhibitory effect in a warm-blooded  
10 animal, such as man.

According to another feature of the invention there is provided the use of a compound selected from Group B or Group D or the Reference Examples, or a pharmaceutically acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an  $11\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man.

- 15        Where production of or producing an  $11\beta$ HSD1 inhibitory effect is referred to suitably this refers to the treatment of metabolic syndrome. Alternatively, where production of an  $11\beta$ HSD1 inhibitory effect is referred to this refers to the treatment of diabetes, obesity, hyperlipidaemia, hyperglycaemia, hyperinsulinemia or hypertension, particularly diabetes and obesity. Alternatively, where production of an  $11\beta$ HSD1 inhibitory effect is referred to this  
20 refers to the treatment of glaucoma, osteoporosis, tuberculosis, dementia, cognitive disorders or depression.

- According to a further feature of this aspect of the invention there is provided a method for producing an  $11\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount  
25 of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

- According to a further feature of this aspect of the invention there is provided a method for producing an  $11\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih) or (II) or a  
30 pharmaceutically acceptable salt thereof, or a compound selected from Group A, Group C or the Examples, or a pharmaceutically acceptable salt thereof.

According to a further feature of this aspect of the invention there is provided a method for producing an  $11\beta$ HSD1 inhibitory effect in a warm-blooded animal, such as man,

in need of such treatment which comprises administering to said animal an effective amount of a compound selected from Group B or Group D or the Reference Examples, or a pharmaceutically acceptable salt thereof.

In addition to their use in therapeutic medicine, the compounds of formula (I), or a  
5 pharmaceutically acceptable salt thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of 11 $\beta$ HSD1 in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

The inhibition of 11 $\beta$ HSD1 described herein may be applied as a sole therapy or may  
10 involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example agents than might be co-administered with 11 $\beta$ HSD1 inhibitors, particularly those of the present  
15 invention, may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide) and prandial glucose regulators (for example repaglinide, nateglinide);
- 3) Insulin sensitising agents including PPAR $\gamma$  agonists (for example pioglitazone and  
20 rosiglitazone);
- 4) Agents that suppress hepatic glucose output (for example metformin);
- 5) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 6) Agents designed to treat the complications of prolonged hyperglycaemia; e.g. aldose  
25 reductase inhibitors
- 7) Other anti-diabetic agents including phosphotyrosine phosphatase inhibitors, glucose 6 - phosphatase inhibitors, glucagon receptor antagonists, glucokinase activators, glycogen phosphorylase inhibitors, fructose 1,6 biphosphatase inhibitors, glutamine:fructose -6-phosphate amidotransferase inhibitors
- 30 8) Anti-obesity agents (for example sibutramine and orlistat);
- 9) Anti- dyslipidaemia agents such as, HMG-CoA reductase inhibitors (statins, eg pravastatin); PPAR $\alpha$  agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors);

- 53 -

- ileal bile acid absorption inhibitors (IBATI), cholesterol ester transfer protein inhibitors and nicotinic acid and analogues (niacin and slow release formulations);
- 10) Antihypertensive agents such as,  $\beta$  blockers (eg atenolol, ideral); ACE inhibitors (eg lisinopril); calcium antagonists (eg. nifedipine); angiotensin receptor antagonists (eg candesartan),  $\alpha$  antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 5 11) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors; antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin; and
- 10 12) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone).

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

15 **Examples**

The invention will now be illustrated in the following non limiting Examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation *in vacuo* and work up procedures were
- 20 carried out after removal of residual solids such as drying agents by filtration;
- (ii) all reactions were carried out under an inert atmosphere at ambient temperature, typically in the range 18-25°C, with solvents of HPLC grade under anhydrous conditions, unless otherwise stated;
- (iii) column chromatography (by the flash procedure) was performed on Silica gel 40-63  $\mu$ m
- 25 (Merck);
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; magnetic resonance chemical shift values were measured in deuterated  $CDCl_3$  (unless otherwise stated)
- 30 on the delta scale (ppm downfield from tetramethylsilane); proton data is quoted unless otherwise stated; spectra were recorded on a Varian Mercury-300 MHz, Varian Unity plus-400 MHz, Varian Unity plus-600 MHz or on Varian Inova-500 MHz spectrometer unless otherwise stated data was recorded at 400MHz; and peak multiplicities are shown as follows:

100804-2

- 54 -

s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; ABq, AB quartet; ABd, AB doublet, ABdd, AB doublet of doublets; dABq, doublet of AB quartets; LCMS were recorded on a Waters ZMD, LC column xTerra MS C<sub>8</sub>(Waters), detection with a HP 1100 MS-detector diode array equipped; mass spectra  
5 (MS) (loop) were recorded on VG Platform II (Fisons Instruments) with a HP-1100 MS-detector diode array equipped; unless otherwise stated the mass ion quoted is (MH<sup>+</sup>); unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on Prep LC 2000 (Waters), Cromasil C<sub>8</sub>, 7 µm, (Akzo Nobel); MeCN and de-ionised water 10 mM ammonium acetate as mobile phases, with  
10 suitable composition;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;

(viii) where solutions were dried sodium sulphate was the drying agent;

(ix) where an "ISOLUTE" column is referred to, this means a column containing 2 g of silica,  
15 the silica being contained in a 6 ml disposable syringe and supported by a porous disc of 54Å pore size, obtained from International Sorbent Technology under the name "ISOLUTE"; "ISOLUTE" is a registered trade mark;

(x) the following abbreviations may be used hereinbefore or hereinafter:-

DCM	dichloromethane;
20 EtOAc	ethyl acetate;
DMSO	dimethylsulphoxide;
DMF	dimethylformamide;
ether	diethyl ether;
LDA	lithium diisopropylamine;
25 MeCN	acetonitrile; and
THF	tetrahydrofuran.

### Example 1

#### (Thien-3-ylmethyl)-(4-chlorophenyl)-ketone

30 A solution of 4-chlorophenyl magnesium bromide in ether (6.0ml of a 1.0 mol solution, 6.0 mmol) was added to a stirred solution of *N*-methoxy-*N*-methyl-3-thienylmethanamide (Method 1; 370 mg, 2.0 mmol) in THF (20 ml) at 0°C. The resultant mixture was stirred at ambient temperature overnight, and then quenched with ethanol (50

100804-2

- 55 -

ml). The resultant mixture was evaporated to dryness and the residue partitioned between water (50 ml) and ether (100 ml). The ether layer was separated, washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 5% EtOAc in hexane as eluent to give the title compound, as a solid (170 mg, 0.72 mmol). NMR: 4.3 (s, 2H), 7.0 (d, 1H), 7.1 (d, 1H), 7.3 (dd, 1H), 7.4 (d, 2H), 7.9 (d, 2H).

### Examples 2-3 and Reference Example 1

The procedure described in Example 1 was repeated using the appropriate Grignard reagent and the appropriate Weinreb derivative to obtain the compounds described below.

Ex	Compound	NMR / m/z
2	(Thien-3-ylmethyl)-(4-fluorophenyl)-ketone	4.3 (s, 2H), 7.0 (d, 1H), 7.1 (m, 3H), 7.3 (dd, 1H), 8.0 (m, 2H)
3	(Thien-3-ylmethyl)-(4-methylphenyl)-ketone	2.4 (s, 3H), 4.3 (s, 2H), 7.0 (d, 1H), 7.1 (d, 1H), 7.3 (m, 3H), 7.9 (dd, 2H)
RE 1 <sup>1</sup>	(3-Bromophenyl)-(4-methoxybenzyl)-ketone	305

10 <sup>1</sup> This compound was synthesised via p-methoxybenzylchloride Grignard and reacted with a Weinreb amide made from 4-bromobenzoic acid. The Grignard reagent was made with the complex [Mg (anthracene)(THF)<sub>3</sub>].

### Reference Example 2

#### 15 (4-Methoxybenzyl)-(4-fluorophenyl)-ketone

Diisopropylamine (7.4ml, 52.8mmol) was added to anhydrous THF under argon. The solution was stirred and cooled to -35°C in a dry ice/acetone bath and *N*-butyl lithium (1.6M, 31.2ml, 50.4mmol) was added via a syringe over 2 – 3 mins, controlling the exotherm to below -20°C. After the addition was complete the reaction was cooled to -70°C and a solution  
 20 of 4-methoxyphenylacetic acid (3.89g, 24mmol) in THF (48ml) was transferred to the reaction mixture via a cannula, adding the solution dropwise and keeping the temperature below -55°C. The reaction was stirred for 10 mins at -70°C and then allowed to warm up to -15°C. A solution of *N*-methyl-*N*-methoxy-4-fluorophenylcarbamoyl (4.38g, 26.4mmol) in THF (48ml) was added via a dropping funnel dropwise, while the reaction mixture  
 25 exothermed to 0°C. After addition was complete, the reaction was allowed to warm up to room temperature and stirred for 1.5 hours. The reaction was added to a stirred solution of

100804-2

- 56 -

concentrated hydrochloric acid (13ml) in water (250ml), and was stirred for 10 mins before the addition of ether. The organic layer was separated, washed with water, aqueous sodium bicarbonate and brine and dried ( $\text{MgSO}_4$ ). The solvent removed *in vacuo* to give a sticky solid. This was triturated with methanol, and the resultant solid was dried under high vacuum to give a white solid (2.2g). Mp 107 - 109°C; NMR(200MHz) 3.78 (3H, s), 4.17 (2H, s), 6.87 (2H, d), 7.13 (4H, m), 8.03 (2H, dd).

### Reference Example 3

#### (4-Bromophenyl)-(4-chlorobenzyl)-ketone

4-Chlorophenylacetic acid (12.78g) was heated with phosphorus trichloride for 1 hour and then bromobenzene (42.5ml) was added. The top layer was decanted onto aluminium chloride (11.25g) in carbon disulfide (50ml) and the bottom layer was washed with additional bromobenzene (42.5ml) before this was decanted as well. The reaction mixture was allowed to stir at this temperature for 1.5 hours before heating on a steam bath for 2 hours. The reaction was cooled, poured onto ice/concentrated hydrochloric acid (300ml) and stirred for 1 hour before extracting with chloroform (3 x 200ml). The organic layer was washed with sodium hydroxide (2 x 200ml) and water (2 x 200ml) and then dried ( $\text{MgSO}_4$ ). The chloroform and any remaining bromobenzene was removed *in vacuo*. On cooling the resultant black oil crystallised. This solid was recrystallized twice from chloroform/petrol to give 12.4g, 54%. Mp 128 - 129°C;  $m/z$  308 ( $\text{M}^+$ ).

### Reference Example 3

The procedure described in Reference Example 3 was repeated using the appropriate starting materials to obtain the compound below.

Ex	Compound	Data
RE 3	(4-Bromobenzyl)-(4-chlorophenyl)-ketone	Mp 125 - 126°C; 308 ( $\text{M}^+$ )

### Example 4

#### (4-Chlorophenyl)-(3,4-dichlorophenylmethyl)-ketone

Magnesium turnings (1.95g) and a crystal of iodine were placed in a reaction vessel and 3,4-dichlorobenzyl chloride (14.63g) in ether (25ml) was added slowly ensuring that the temperature of the reaction mixture was kept at the reflux temperature of ether. After the addition was complete the reaction was refluxed for a further 30 mins. 4-Chlorobenzonitrile

100804-2

- 57 -

(8.25g) in ether (50ml) was added dropwise while the reaction was at a temperature of  $\sim 30^{\circ}\text{C}$ . After the addition was complete the reaction was refluxed for 3 hours before being poured onto ice/concentrated sulphuric acid (1 litre) and left to stand overnight. The solid that remained was filtered, washed with water, dissolved in DCM and dried ( $\text{MgSO}_4$ ), filtered and 5 evaporated to dryness. The resulting solid was recrystallized from DCM and petrol twice. (6.05g, 46%). Mp  $99 - 103^{\circ}\text{C}$ ;  $m/z$  298 ( $\text{M}^+$ ).

**Example 5**

The procedure described in Example 4 was repeated with the appropriate starting 10 materials to obtain the compound described below.

Ex	Compound	Mp / Mz
5	(3-Bromophenyl)-(4-chlorophenylmethyl)-ketone	$64 - 67^{\circ}\text{C}$ ; 308 ( $\text{M}^+$ )

**Example 6****(4-Bromo-2-hydroxyphenyl)-(4-bromobenzyl)-ketone**

(3-Bromophenoxy)-(4-bromobenzyl)-ketone (Method 3; 36.5g) and aluminium 15 trichloride (26.3g) were dissolved in nitrobenzene (100ml) and the mixture was stirred at  $100^{\circ}\text{C}$  for 2.5 hours. The reaction was left to stand overnight. The reaction mixture was poured onto a mixture of ice/concentrated hydrochloric acid and stirred for 5 mins. The aqueous layer was extracted with EtOAc. The organic layers were combined and the solvent removed *in vacuo*, and the nitrobenzene was removed by steam distillation. The resultant 20 residue was purified by column chromatography to give the product 31g. NMR ( $\text{DMSO}-d_6$ ; 400MHz) 4.30 (s, 2H), 7.05 (dd, 1H), 7.25 (m, 3H), 7.40 (m, 2H), 7.80 (d, 1H), 11.70 (bs, 1H);  $m/z$  370.

**Reference Example 5****25 (4-Methoxyphenyl)-(benzyl)-ketone**

Sodium hydride (50% dispersion in oil, 960mg, 20mmol) was washed with petrol and suspended in anhydrous DMF (10ml) under nitrogen. *N,N*-Diethyl-*N*-( $\alpha$ -cyano-4-methoxybenzyl)amino (Method 4: 3.27g, 15mmol) was placed in anhydrous DMF (20ml) and added to the reaction which was stirred at room temperature for 1 hour. Benzyl chloride 30 (1.89g, 1.73ml, 15mmol) in anhydrous DMF (10ml) was added dropwise over 1 hour at room temperature (slight exotherm) and the reaction was stirred overnight at room temperature.



100804-2

- 58 -

Methanol (5ml) was added and the solvent removed *in vacuo* at 90°C followed by high vacuum for 2 hours, to give a yellow oil. This was stirred in hydrochloric acid (6M; 40ml) for 16 hours, and then extracted into chloroform (3 x 30ml), washed with water, dried (MgSO<sub>4</sub>) and evaporated to dryness to give a yellow oil. This was purified by column chromatography with ether:hexane (2:1). The product was then recrystallized from 40 - 60°C petrol to give a colourless solid 410mg. Mp 68 - 69°C; m/z 226.

### Reference Example 6

#### (4-Methylthiobenzyl)-(4-fluorophenyl)-ketone

- 10 To 10ml of sieve-dried DCM containing anhydrous zinc iodide (50mg) was added 4-fluorobenzaldehyde (630mg, 0.54ml, 5mmol). To this stirred mixture, at room temperature and under argon, was added trimethylsilyl cyanide (520mg, 0.7ml, 5.25mmol) and the reaction was stirred overnight (~20 hours). The solvent was evaporated *in vacuo* and the residue was treated with anhydrous ether (15ml) and a little magnesium sulphate. The solution
- 15 was filtered and the solvent removed *in vacuo* to give the cyanohydrin as an orange oil (1.14g). Lithium diisopropylamide was made from *n*-butyl lithium (2.5M in hexanes, 2.1ml, 5.25mmol) and diisopropylamine (500mg, 0.69ml, 5mmol) in THF (5ml) at -40°C. The reaction was then cooled to -60°C and the cyanohydrin in THF (5ml) was added, under argon, at such a rate as to keep the temperature below -55°C. After this addition the reaction was
- 20 stirred for 30mins and then 4-methylthiobenzyl chloride (900mg, 5.25mmol) was added in THF (2.5ml). The cooling bath was removed and the reaction stood at room temperature overnight. To the reaction was added saturated ammonium chloride solution (13ml) and ether (25ml). The organic phase was separated, washed with saturated ammonium chloride solution, dried (MgSO<sub>4</sub>) and the solvent was removed *in vacuo* to give an orange oil (1.78g).
- 25 The oil was taken up in methanol (7ml) and treated with 2M sulphuric acid (10ml), the oil precipitated and so acetone (20ml) was added to give a clear solution. This stood at room temperature overnight. The pH was adjusted to 7.5 with 2M sodium hydroxide solution and the solution was concentrated *in vacuo*. To the residue was added water and this was extracted with DCM (2x 30ml). The combined extracts were washed with brine and evaporated to give
- 30 a sticky solid (1.3g). This was purified by MPLC (1:5 EtOAc:hexane) to give an off-white flaky solid (930mg). NMR (DMSO-d<sub>6</sub>; 400MHz): 2.40 (s, 3H), 4.30 (s, 2H), 7.20 (s, 4H), 7.35 (tt, 2H), 8.10 (m, 2H); m/z 260.

100804-2

- 59 -

**Reference Example 7****(Pyrid-4-ylmethyl)-(pyrid-4-yl)-ketone**

A solution of methyl lithium (1.4M) in ether was stirred at room temperature under an argon atmosphere, and 4-picoline (3.72g, 3.9ml, 40mmol) was added dropwise over five mins. When the addition was complete the solution was refluxed gently for 30mins and then a solution of methyl isonicotinate (2.75ml, 20mmol) in ether (5ml) was added. The resulting sticky suspension was refluxed for 30 mins, cooled and then water (7.5ml) was added carefully. The reaction mixture was then treated with a cold mixture of 6M hydrochloric acid (10ml) and water (50ml). The ether phase was extracted several times with 6M hydrochloric acid, and the combined aqueous phases were treated with 70% sodium hydroxide solution until the solution remained slightly acid. Solid sodium hydrogen carbonate was then added until the mixture was slightly basic (pH 8). The basic mixture was extracted with ether until the extracts no longer gave a brown colour with alcoholic ferric chloride. The combined ether extracts were dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to give a yellow solid. This was purified by MPLC (7.5% methanol in DCM) to give a yellow solid (2.0g). This was recrystallized from toluene/hexane. NMR: 4.35 (s, 2H), 7.20-7.25 (m, 2H), 7.77-7.82 (m, 2H), 8.60-8.65 (m, 2H), 8.85-8.90 (m, 2H);  $m/z$  199.

**Reference Example 8****(Pyrid-2-ylmethyl)-(phenyl)-ketone**

A solution of *n*-butyl lithium (40ml, 1.6M in hexanes, 66mmol) in anhydrous ether (50ml) was stirred in an argon atmosphere. 2-Picoline (6.5ml, 66mmol) was added over approximately 10mins. The resulting red solution was heated gently for 30 mins then cooled to room temperature. To this rapidly stirred solution of 2-picolyllithium was added a solution of benzonitrile (6.8ml, 66mmol) in anhydrous ether (10ml) dropwise, which gave an orange suspension almost immediately. The reaction was stirred at room temperature for 2 hours. The reaction mixture was then treated with water (50ml) followed by 2M sulphuric acid (50ml) and the two phase mixture was heated under reflux for 30 mins. After cooling, the reaction mixture was extracted with ether. The aqueous solution was adjusted to pH 7, and then extracted further with ether. The combined ether solutions were washed with water, dried ( $\text{MgSO}_4$ ), and the solvent removed *in vacuo* to give a deep yellow oil. This was purified by column chromatography (60 – 80°C petroleum ether/EtOAc 2:1) and the resulting yellow oil crystallised from warm 60 – 80°C petroleum ether to give long yellow needles (410mg). Mp

100804-2

- 60 -

57 – 59°C; NMR: 3/2 mixture of keto/enol NOTE first two signals are in a ratio for  $\text{CH}_2\text{C}(\text{O})$  and  $\text{CH}=\text{CHOH}$  4.50 (s), 6.05 (s) 8.60-6.90 (m, 9H), OH offset.

#### Reference Example 9

##### 5 (Benzyl)-[4-(tetrahydropyran-2-yloxy)phenyl]-ketone

4-Hydroxydeoxybenzoin (20g) was placed in dihydropyran, with 2 drops of concentrated hydrochloric acid. The mixture was heated for 4 hours at 55°C, and then cooled. The resultant precipitate was taken up in ether:toluene (1:1), heated until the solid went into solution, washed with aqueous sodium hydroxide (2 x 50ml), water (2 x 50ml), dried, and the  
10 solvent removed *in vacuo* to give a yellow solid. This was taken up in the minimum amount of ethanol with charcoal, and heated. The solution was filtered and as the solution cooled, the product crystallised out. This was separated by filtration and dried in a dessicator. (22.6g).  
NMR ( $\text{DMSO}-d_6$ ; 400MHz): 1.50 – 1.80 (m, 3H), 1.80 – 1.95 (m, 3H), 3.55 – 3.60 (m, 1H), 3.75 – 3.85 (m, 1H), 4.30 (s, 2H), 5.60 (m, 1H), 7.10 (m, 2H), 7.20 – 7.35 (m, 5H), 8.00 (m,  
15 2H);  $m/z$  297.

#### Reference Example 10

##### [4-(Benzyl)morpholin-2-ylmethyl]-(phenyl)-ketone

Lithium (2.8g) was cut into small pieces under an atmosphere of argon, and placed  
20 into sodium dried ether (100ml). A small portion of bromobenzene dissolved in ether was added with vigorous stirring – a reaction commenced within 5 mins and the remaining bromobenzene (21.0ml in 100ml of sodium-dried ether) was added dropwise over 30mins, maintaining a gentle reflux. The reaction was refluxed for a further hour, and then cooled to -20°C. (4-Benzylmorpholin-2-yl)acetonitrile (Journal of Medicinal Chemistry (1990), 33(5),  
25 1406-1321.6g) was dissolved in sodium dried ether (108ml) and added dropwise over 15 mins, the temperature not rising above -15°C. The reaction was stirred at this temperature for 15 mins, and then the reaction mixture was poured into 2M hydrochloric acid (800ml) and ice water (800ml) with stirring. This was stirred at room temperature for 15 mins, the ether later separated, and the aqueous layer was separated and washed with ether. The aqueous layer was  
30 carefully basified with sodium carbonate and extracted with ether. The combined ether extracts were washed with brine, dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to give the product as a light brown oil, and as the hydrochloric acid salt. (24.9g). NMR ( $\text{DMSO}-d_6$ ;

100804-2

- 61 -

400MHz) 2.90 – 3.15 (m, 2H), 3.15 – 3.50 (m, 2H), 3.80 – 4.05 (m, 2H), 4.30 (s, 3H), 7.35 – 7.80 (m, 8H), 7.95 (d, 2H), 11.00 (bs, 1H); m/z 295.

### Example 7

#### 5 (Benzimidazol-1-ylmethyl)-(2,4-dichlorophenyl)-ketone

Benzimidazole (5.9g) was added to a solution of sodium hydride (2.4g, 50% dispersion in oil) in DMF (55ml) and stirring was continued until effervescence ceased (25mins). To this brown solution was added 2, 2',4'-trichloroacetophenone (11.17g) in DMF (35ml) over 15mins and the resulting brown solution was stirred at room temperature for 2  
10 hours. The reaction mixture was poured into water and this was extracted with EtOAc. The extracts were washed with water, dried and the solvent removed *in vacuo* to give a dark red oil. This was purified by column chromatography (chloroform:MeOH:NH<sub>3</sub> 9:1:0.1) to give an orange oil. This was purified by column chromatography (EtOAc) to give a pale yellow oil (2.5g). Mp 130 – 132°C; NMR (DMSO-d<sub>6</sub>; 400MHz): 5.90 (s, 2H), 7.20 – 7.30 (m, 2H), 7.55  
15 (m, 1H), 7.65 – 7.70 (m, 2H), 7.85 (d, 1H), 8.10 (d, 1H), 8.20 (s, 1H); m/z 305.

### Example 8

#### [1-Methyl-1-(1,2,4-triazol-1-yl)ethyl]-(4-trifluoromethyl-2-fluorophenyl)-ketone

To (2-fluoro-4-trifluoromethylphenyl)-(2-bromoprop-2-yl)-ketone (Method 5; 9.0g)  
20 was added sodium triazole (2.9g) in DMF (50ml). The mixture was heated at 70°C for 1.5 hours, and the solvent removed. The resulting mixture was purified by MPLC (DCM graduating to 5% methanol in DCM) followed by recrystallization from EtOAc/hexane. NMR (DMSO-d<sub>6</sub>; 400MHz): 1.90 (s, 6H), 7.25 (m, 1H), 7.55 (m, 1H), 7.75 (d, 1H), 8.75 (s, 1H); m/z 302.

25

### Reference Example 11

#### (Benzyl)-(4-methylphenyl)-ketone

A solution of phenylacetyl chloride (77.3g) in toluene (250ml) was added dropwise to a suspension of aluminium trichloride (80g) in toluene (150ml) over 30mins with the  
30 temperature of the reaction not exceeding 60°C. The reaction was stirred at room temperature for 2.5 hours, then heated at 60°C for a further 1.5 hours. The reaction mixture was cooled and poured onto ice/hydrochloric acid. The layers were separated and the aqueous layer extracted with toluene. The organic layers were combined, dried, and the solvent removed *in vacuo*.

100804-2

- 62 -

The solid was recrystallized from 80 – 100°C petroleum ether, and dried by heating to give a yield of 77.7g. Mp 107 – 111°C; NMR (DMSO-d<sub>6</sub>, 400MHz): 2.35 (s, 3H), 4.30 (s, 2H), 7.15 – 7.35 (m, 7H), 7.95 (d, 2H); m/z 211.

## 5 Reference Example 12

### (Imidazol-1-ylmethyl)-(2-chlorothiophen-5-yl)-ketone

To a solution of 2-chloro-5-acetylthiophene (Method 8; 32g) in chloroform (250ml) was added bromine (32g, 10ml) in chloroform (100ml) over a period of 1.5 hours. The reaction was catalysed by a few drops of HBr/AcOH and by a UV lamp which also kept the temperature at 40 – 45°C during the addition. After the addition was complete, the stirring was continued at room temperature for 2 hours. The reaction mixture was poured onto water and the organic layer was separated and washed with water, dried and the solvent removed *in vacuo* to give a brown oil which solidified on standing (41.4g). The solid was dissolved in DMF (100ml) and added dropwise to a stirred solution of imidazole (68g) in DMF (200ml) at 5 – 10°C, over 30 mins. The resultant brown solution was stirred at room temperature for 18 h. The reaction mixture was poured onto water and extracted with EtOAc. The extracts were washed with water, dried and evaporated to a black gum which crystallised on standing. This solid was re-crystallised from EtOAc to give a tan coloured solid which was filtered, and washed with ether to give 11.5g. Mp 109 – 111°C; NMR (DMSO-d<sub>6</sub>): 5.18 (s, 2H), 6.65 – 7.10 (m, 3H), 7.22 – 7.48 (m, 2H).

## Example 9

### [2-(4-Chlorophenyl)-1-(pyrid-3-yl)ethyl]-(4-chlorophenyl)-ketone

Sodium hydride (100%) (500mg) was suspended in anhydrous DMF (30ml) and the reaction was cooled to 0°C. To the reaction was added (pyrid-3-ylmethyl)-(4-chlorophenyl)-ketone (Reference Example 13) in DMF (20ml) and the reaction mixture was stirred for 1 hour. 4-Chlorobenzyl chloride (3.2g) in DMF (10ml) was added, and the reaction stirred at 0°C for 3 hours. The reaction was poured onto water, and the resultant crystals were filtered and recrystallized from 60 – 80°C petrol with charcoal to give a white solid (2.75g). Mp 98 – 99°C.

100804-2

- 63 -

**Example 10****[ $\alpha$ -(4-Fluorobenzyl)benzyl]-(pyrid-3-yl)-ketone**

The procedure described in Example 9 was repeated substituting the 4-chlorobenzyl chloride with 4-fluorobenzyl chloride and using (pyrid-3-yl)-(benzyl)-ketone (Reference Example 14) to give the title compound 7g. Mp 100 - 102°C; m/z 305 (M<sup>+</sup>).

**Reference Example 13****(Pyrid-3-ylmethyl)-(4-chlorophenyl)-ketone**

To diisopropylamine (56ml) was added *n*-butyl lithium (2.4M, 166ml) keeping the temperature of the reaction mixture below 20°C by cooling in an ice/salt bath. 3-Picoline (37.2g) was added dropwise, diluted with toluene (30ml) and the reaction stirred at 0°C for 30 mins before the addition of 4-chloromethyl benzoate (34g) in anhydrous toluene (30ml). The reaction was stirred at 5°C for 1.5 hours. The reaction mixture was poured onto ice/water, acidified and washed to remove ester, then the acid layer was basified and extracted to give a red oil. This was distilled under vacuum which gave 1 fraction at 154°C, 0.2mmHg, which resulted in yellow crystals. These were triturated with petrol/ether to give a cream solid (17g). Mp 61 - 63°C.

**Reference Example 14****(Pyrid-3-yl)-(benzyl)-ketone**

The title compound was prepared by the procedure of Reference Example 13 using the appropriate starting materials. Mp 132-138°C.

**Example 11****(4-Chlorophenyl)-{ $\alpha$ -hydroxy- $\alpha$ -[1-(1,2,4-triazol-1-yl)ethyl]-4-chlorobenzyl}-ketone**

A solution of 4-bromo-1-chlorobenzene (17.1g, 90mmol) in anhydrous ether (100ml) was added portionwise to magnesium turnings (2.16g, 90mmol) suspended in anhydrous ether (100ml), with a crystal of iodine. The reaction was refluxed for 4 hours. Anhydrous toluene (150ml) was added and the temperature increased, with the ether distilling off. When all of the ether had been removed, 1-[1-(4-chlorophenyl)-1-(trimethylsilyloxy)-1-(cyano)prop-2-yl]-1,2,4-triazole (Method 9) in toluene (10ml) was added and the reaction was refluxed and stirred overnight. After cooling, the reaction mixture was acidified with 3M hydrochloric acid (100ml) and stirred for 1 hour. The aqueous and organic layers were separated, and the

100804-2

- 64 -

aqueous layers washed with ether. The ether extracts were combined with the organic layer, which was then dried ( $\text{MgSO}_4$ ). After filtration, the solvent was removed *in vacuo* to give a deep red oil. The was purified by column chromatography (4% methanol in DCM) to give the required product (1.4g). Mp 181 – 183°C; NMR: 1.70 (d, 3H), 5.30 (q, 1H), 5.70 (s, 1H), 7.50 (m, 10H);  $m/z$  376.

#### Reference Example 15

##### [2,4-Dichloro- $\alpha$ -(1,2,4-triazolylmethyl)benzyl]-(4-chlorophenyl)-ketone

[1-(2,4-Dichlorophenyl)vinyl]-(4-chlorophenyl)-ketone (Method 24; 1.6g, 5mmol)  
10 was added to ethanol (25ml) containing triazole (2g) and triethylamine (20 drops) and the reaction was stirred at room temperature for 2 hours. The reaction mixture was poured onto water and extracted with ether. The extracts were washed with water, and the solvent evaporated *in vacuo* to give an oil. This was placed in ether/petrol to give a white precipitate which was collected by filtration. This was purified by MPLC to give a free flowing white  
15 crystalline solid (1.25g). Mp 109 - 111°C; NMR: 4.35 (q, 1H), 4.95 (q, 1H), 5.65 (q, 1H), 7.20 (m, 5H), 7.75 (m, 2H), 7.85 – 8.00 (dd, 2H).

#### Reference Example 16

##### (4-Chlorophenyl)-[ $\alpha$ -(1,2,4-triazol-1-ylmethyl)-4-chlorobenzyl]-ketone

20 Reference Example 15 was repeated with [1-(4-chlorophenyl)vinyl]-(4-chlorophenyl)-ketone (J. Med. Chem. (1972), 15(12), 1243-7) to give the title compound. Mp 126 - 128°C. 345 ( $M^+$ ).

#### Reference Example 17

##### (2,4-Dichlorobenzyl)-(4-chlorophenyl)-ketone

2,4-Dichlorobenzyl chloride (92.5g, 0.48mol) in ether (300ml) was added over 1 hour to magnesium (13g) in ether (50ml) at reflux, and the reaction was allowed to stand at room temperature overnight. 4-Chlorobenzonitrile (0.2mol) was dissolved in sieve-dried THF and the Grignard reagent (180ml) was added over 5 min with stirring. The reaction mixture was  
30 refluxed for 24 h under argon. The reaction mixture was cooled and poured into 2M hydrochloric acid/ice and extracted with EtOAc. On drying of the solution and evaporation of the solvent gave a yellow solid. This was triturated with 50:50 EtOAc:ether, and the resulting pale yellow solid was filtered (24.7g). Mp 127 - 129°C.

100804-2

- 65 -

**Example 12****[2-(2-Fluorophenyl)-1-(1,2,4-triazol-1-yl)ethyl]-(thien-2-yl)-ketone**

Sodium hydride (610mg, 26mmol) was suspended in DMF (10ml) and (1,2,4-triazol-1-ylmethyl)-(thien-2-yl)-ketone (Journal of Medicinal Chemistry (1987), 30(8), 1497-502; 5g, 26mmol) dissolved in DMF (30ml), was added. The reaction was stirred at room temperature for 3 hours and then cooled in an ice-bath and 2-fluorobenzyl chloride (3.72g, 26mmol), in DMF (15ml) was added dropwise, keeping the temperature between 0 and 5°C. The reaction was left to stir at room temperature overnight, then poured onto water which formed a precipitate. This was filtered and recrystallized from petroleum ether 60 - 80°C to give the product (2.95g). Mp 121 - 122°C.

**Example 13****[2-(4-Chlorophenyl)-1-(pyridazin-3-yl)ethyl]-(phenyl)-ketone**

(Phenyl)-(pyridazin-3-ylmethyl)-ketone (Chemical & Pharmaceutical Bulletin (1978), 26(12), 3633-40.2.5g, 13mmol) in DMF (25ml) was added to a suspension of sodium hydride (610mg, 50% dispersion in oil, 13mmol, washed with ether) in DMF (10ml). After stirring for 2 hours the solution was cooled in an ice/salt bath and 4-chlorobenzyl chloride (2g, 12.5mmol) in DMF (15ml) was added dropwise at 0 - 5°C. The reaction mixture was warmed to room temperature and stirred for a further 1 hour. The reaction mixture was poured onto water (200ml) which gave a yellow precipitate, which was filtered, washed with water, dried and recrystallized from EtOAc/60 - 80°C petroleum ether to give the product (1.6g). Mp 140 - 142°C; m/z 322 (M<sup>+</sup>).

**Examples 14-16**

Following the procedure described in Example 13, the following compounds were made using the appropriate starting materials.

Ex	Compound	Data
14 <sup>1</sup>	[2-(2,4-Dichlorophenyl)-1-(pyridazin-3-yl)ethyl]-(phenyl)-ketone	NMR (400MHz; DMSO-d <sub>6</sub> ): 3.35 (m, 1H), 3.55 (m, 1H), 5.50 (m, 1H), 7.30 (s, 2H), 7.45 (m, 2H), 7.50 - 7.70 (m, 3H), 7.95 (d, 2H), 9.05 (m, 1H); m/z 359
15 <sup>2</sup>	[2-(4-Chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyridin-3-yl)-ketone	Mp 116 - 118°C



100804-2

- 66 -

16 <sup>2</sup>	[2-(4-Chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(thien-2-yl)-ketone	Mp 107 - 109°C; m/z 312 (M <sup>+</sup> )
-----------------	--	---

<sup>1</sup> Prep of starting material: Chemical & Pharmaceutical Bulletin (1978), 26(12), 3633-40

<sup>2</sup> The starting materials for Examples 15 and 16 could be prepared according to the procedure described in Chemical & Pharmaceutical Bulletin (1978), 26(12) for (phenyl)-(pyridazin-3-ylmethyl)-ketone.

5

### Examples 17 and 18

[2-(4-Chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyridin-3-yl)-ketone enantiomer 1 and [2-(4-Chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyridin-3-yl)-ketone enantiomer 2

[2-(4-Chlorophenyl)-1-(pyrazin-2-yl)ethyl]-(pyridin-3-yl)-ketone (Example 15) was

10 separated into its 2 enantiomers using the following HPLC conditions.

Instrument	Perkin Elmer 200
Column	10µm Chiralpak AD (4.6mm x 250mm) No. AD00CE-BJ182
Eluent	MeCN/MeOH 95/5
Oven Temperature	Ambient
Flow	1ml/min
Wavelength	254nm
Sample Concentration	1mg/ml in EtOH
Sample Volume	20µl
Run Time	30mins

### Reference Example 18

#### (Phenyl)-(cyclohexylmethyl)-ketone

In a conical flask was placed deoxybenzoin (500mg, 2.55mmol), tetrabutylammonium  
 15 bromide (41mg, 0.13mmol), (bromomethyl)cyclohexane (1.35g, 7.65mmol), toluene (18ml)  
 and 45% KOH in water (6ml). The reaction was sonicated at room temperature for 3 hours  
 and then quenched with saturated ammonium chloride solution (~5ml). The volatiles were  
 removed under reduced pressure and the resulting material was partitioned between ether and  
 water. The organic layer was separated and re-extracted with water then washed with brine,  
 20 dried (MgSO<sub>4</sub>), filtered and evaporated to yield an oil which was further purified by prep

100804-2

- 67 -

LCMS to yield the product as a clear oil. NMR: 1.00 (br m, 2H), 1.25 (br m, 3H), 1.70 (br m, 5H), 2.00 (m, 1H), 2.80 (d, 2H), 7.45 (t, 2H), 7.55 (t, 1H), 7.95 (d, 2H); m/z: 202.

**Example 19****5 (4-Fluorophenethyl)-(4-trifluoromethylphenyl)-ketone**

[2-(4-Fluorophenyl)vinyl]-(4-trifluoromethylphenyl)-ketone (Method 25; 1g) was hydrogenated over Pd/CaCO<sub>3</sub> in ethanol. The catalyst was filtered off, the solvent removed *in vacuo* and the residue obtained recrystallized from aqueous ethanol (510mg). Mp 66 - 67°C; m/z 296 (M<sup>+</sup>).

10

**Examples 20-22 and Reference Example 19**

The procedure described in Example 19 was carried out using the appropriate starting materials to obtain the products described below.

Ex	Compound	Data	SM
20	(4-Fluorophenethyl)-(4-chlorophenyl)-ketone	Mp 60°C; m/z 262 (M <sup>+</sup> )	Method 26
21	(4-Chlorophenethyl)-(2,4-difluorophenyl)-ketone	Mp 70 - 71°C; m/z 280 (M <sup>+</sup> )	Method 27
22	(4-Fluorophenethyl)-(2,4-difluorophenyl)-ketone	Mp 46°C; m/z 264 (M <sup>+</sup> )	Method 28
RE 19	(Phenethyl)-(4-methoxyphenyl)-ketone	NMR (400MHz, DMSO-d <sub>6</sub> ): 2.80 (t, 2H), 3.20 (t, 2H), 3.85 (s, 3H), 6.90 (m, 2H), 7.10 (m, 1H), 7.20 (m, 4H), 7.85 (m, 2H); m/z 240 (M <sup>+</sup> )	Method 29

**15 Example 23****(Phenyl)-[2-(4-methylphenyl)-1-(piperidin-1-yl)ethyl]-ketone**

4-Methylbenzylideneacetophenone (11.0g, 50mmol) and piperidine (17ml, 230mmol) were heated in a sealed tube at 100°C for 4 hours. The mixture was cooled to room temperature, the solid product filtered and crystallised from hexane to give the title compound as a solid (7.0g, 23mmol). Mp. 71-72°C; m/z 307 (M<sup>+</sup>).

100804-2

- 68 -

**Example 24****( $\alpha$ -Methylamino-4-methylbenzyl)-(4-methylphenyl)-ketone**

To 4,4'-dimethylbenzoin (500mg, 2.1 mmol) in 40% aq methylamine (1.1ml) was added methylamine hydrochloride (20mg). The reaction was warmed to reflux and stirred at this temperature for 2 hours before addition of further 40% aq methylamine (0.5ml). The reaction was stirred at reflux for a further 3 hours then cooled to room temperature. Saturated sodium hydrogen carbonate (15ml) was added and the crude mixture was extracted with ether (2x30ml). The ether layers were combined and washed with brine then dried (MgSO<sub>4</sub>), filtered and evaporated under reduced pressure to yield an oil. This crude product was dissolved in ether and then acidified with hydrochloric acid in ether (~0.2M), the resulting precipitate was filtered off and recrystallized from EtOH to give the product as a white solid (154mg, 25%). NMR (DMSO-d<sub>6</sub>): 2.25 (s, 3H), 2.35 (s, 3H), 2.45 (s, 3H), 6.35 (s, 1H), 7.25 (d, 2H), 7.30 (d, 2H), 7.45 (d, 2H), 7.90 (d, 2H), 9.90 (br s, 2H); m/z 254.

**15 Examples 25-28**

The procedure described in Example 24 was repeated using the appropriate starting materials to obtain the compounds described below.

Ex	Compound	M/z	NMR (DMSO-d <sub>6</sub> )
25	( $\alpha$ -Methylamino-4-chlorobenzyl)-(4-chlorophenyl)-ketone	294	2.45 (s, 3H), 6.50 (s, 1H), 7.50 (d, 2H), 7.60 (d, 4H), 8.05 (d, 2H), 9.80 (br s, 1H), 10.2 (br s, 1H)
26	( $\alpha$ -Ethylamino-4-chlorobenzyl)-(4-chlorophenyl)-ketone	308	1.30 (t, 3H), 2.80 (br s, 1H), 2.95 (br s, 1H), 6.50 (br s, 1H), 7.55 (d, 2H), 7.60 (m, 4H), 8.10 (d, 2H), 9.65 (br s, 1H), 9.90 (br s, 1H)
27	( $\alpha$ -Isopropylamino-4-chlorobenzyl)-(4-chlorophenyl)-ketone	322	1.30 (m, 6H), 3.05 (br s, 1H), 6.40 (br s, 1H), 7.50 (d, 2H), 7.60 (d, 2H), 7.70 (d, 2H), 8.15 (d, 2H), 9.50 (br s, 1H)
28	( $\alpha$ -Ethylamino-4-methylbenzyl)-(4-methylphenyl)-ketone	268	1.25 (t, 3H), 2.25 (s, 3H), 2.30 (s, 3H), 2.70 (br s, 1H), 2.90 (br s, 1H), 6.35 (s, 1H), 7.20 (d, 2H), 7.30 (d, 2H), 7.45 (d, 2H), 7.95 (d, 2H), 9.50 (br s, 1H), 9.95 (br s, 1H)

100804-2

- 69 -

**Example 29****(1,3-Benzodioxol-5-yl)-[1-(1,3-benzodioxol-5-yl)-1-(ethylamino)methyl]-ketone**

A suspension of piperoin (250mg, 0.83mmol) and ethylamine hydrochloride (40mg, 0.5mmol) in 70% aq ethylamine (4ml) was heated in a microwave at 125°C for 10 minutes. Volatiles were removed under reduced pressure and the resulting crude oil was purified by column chromatography (DCM to 5% MeOH/DCM). This material was dissolved in ether and treated with hydrochloric acid in ether. The resulting solid was filtered off and recrystallized from ethanol to yield a solid (50mg, 20%). NMR (DMSO-d<sub>6</sub>): 1.25 (t, 3H), 2.85 (m, 2H), 6.00 (d, 2H), 6.10 (s, 2H), 6.15 (s, 1H), 6.90 (d, 1H), 7.00 (d, 1H), 7.10 (m, 2H), 7.50 (s, 1H), 7.70 (d, 1H); m/z: 328.

**Example 30****(Thien-2-yl)-[4-(4-chlorobenzoyl)piperidin-1-ylmethyl]-ketone**

To a stirred suspension of (4-chlorophenyl)(4-piperidyl)methanone hydrochloride (100mg, 0.41mmol) in DCM (5ml) was added triethylamine (104mg, 1.03mmol) and 2-bromo-1-(2-thienyl)-1-ethanone (76mg, 0.37mmol). The reaction was stirred at room temperature for 1 hour. The crude reaction mixture was transferred to a separating funnel and washed with 2M hydrochloric acid. The organic layer was separated and washed with water then evaporated to yield an impure solid. This material was partitioned between DCM and saturated sodium hydrogencarbonate solution. The organic layer was separated and washed with brine then dried (MgSO<sub>4</sub>), filtered and evaporated to give a solid. This solid was dissolved in ether and treated with hydrochloric acid in ether. The resulting solid was filtered off to yield the product as a solid (24mg, 17%). NMR (DMSO-d<sub>6</sub>): 2.00 (m, 4H), 3.20 (m, 2H), 3.50 (m, 1H), 3.60 (m, 2H), 5.00 (s, 2H), 7.35 (s, 1H), 7.60 (d, 2H), 8.05 (d, 2H), 8.10 (s, 1H), 8.20 (d, 1H), 10.20 (br s, 1H); m/z: 348.

**Example 31****(α-Methyl-α-hydroxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone**

A solution of methyl magnesium chloride in THF (0.67ml of a 3.0 mol solution, 2.0 mmol) was added to a stirred solution of 4,4'-difluorobenzil (492 mg, 2.0 mmol) in ether (20 ml) during 30 mins at ambient temperature. The resultant mixture was stirred at ambient temperature for 30 mins and then quenched with a saturated aqueous solution of ammonium chloride (2.0 ml) and water (3.0 ml). The ether layer was separated, washed with brine, dried

100804-2

- 70 -

and evaporated to dryness. The residue was purified by column chromatography using 20% EtOAc in hexane as eluent to give the title compound as a solid (300 mg, 1.15 mmol). NMR: 1.9 (s, 3H), 4.5 (s, 1H), 7.0 (m, 4H), 7.4 (dd, 2H), 7.75 (dd, 2H).

#### 5 Examples 32-34 and Reference Example 20

The procedure described in Example 31 was repeated using the appropriate Grignard reagent to replace the methyl magnesium chloride and the appropriate benzil to replace the 4,4'-difluorobenzil to obtain the compounds described below.

Ex	Compound	NMR
32	( $\alpha$ -Benzyl- $\alpha$ -hydroxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone	3.3 (d, 1H), 3.7 (d, 1H), 3.7 (s, 1H), 6.95 (m, 4H), 7.05 (m, 2H), 7.2 (m, 3H), 7.45 (dd, 2H), 7.8 (dd, 2H)
33	( $\alpha$ -Ethyl- $\alpha$ -hydroxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone	0.9 (t, 3H), 2.4 (q, 2H), 4.5 (s, 1H), 7.0 (m, 4H), 7.4 (dd, 2H), 7.7 (dd, 2H)
RE 20	( $\alpha$ -Methyl- $\alpha$ -hydroxy-4-chlorobenzyl)-(4-chlorophenyl)-ketone	1.9 (s, 3H), 4.4 (s, 1H), 7.3 (m, 6H), 7.6 (m, 2H)
34	( $\alpha$ -Methyl- $\alpha$ -hydroxy-2-thienylmethyl)-(2-thienyl)-ketone	2.0 (s, 3H), 4.75 (s, 1H), 7.0 (m, 4H), 7.15 (d, 1H), 7.3 (d, 1H), 7.6 (m, 2H)

#### 10 Example 35

##### ( $\alpha$ -Ethoxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone

A solution of ethyl magnesium bromide in THF (6.0ml of a 1.0 mol solution, 6.0 mmol) was added to a stirred solution of 4,4'-difluorobenzil (492 mg, 2.0 mmol) in ether (20 ml) during 30 minutes at ambient temperature. The resultant mixture was stirred at ambient temperature for 30 minutes and then quenched with a saturated aqueous solution of ammonium chloride (6.0 ml) and water (6.0 ml). The ether layer was separated, washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in hexane as eluent to give the title compound as a solid (58 mg, 0.21 mmol). NMR: 1.2 (t, 3H), 3.6 (q, 4H), 5.4 (s, 1H), 7.0 (m, 4H), 7.4 (dd, 2H), 8.0 (dd, 2H).

100804-2

- 71 -

**Example 36****( $\alpha$ -Isopropoxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone**

A solution of isopropyl magnesium chloride in THF (3.0 ml of a 2.0 mol solution, 6.0 mmol) was added to a stirred solution of 4,4'-difluorobenzil (492 mg, 2.0 mmol) in ether (50 ml) during 30 minutes at ambient temperature. The resultant mixture was stirred at ambient temperature for 30 minutes and then quenched with a saturated aqueous solution of ammonium chloride (6.0 ml) and water (6.0 ml). The ether layer was separated, washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 20% EtOAc in hexane as eluent to give the title compound as a solid (130 mg, 0.45 mmol). NMR: 0.75 (d, 3H), 0.95 (d, 3H), 2.2 (m, 1H), 5.2 (s, 1H), 7.0 (m, 4H), 7.2 (dd, 2H), 7.4 (dd, 2H).

**Example 37****( $\alpha$ -Methoxy-4-fluorobenzyl)-(4-fluorophenyl)-ketone**

Sodium *tert*-butoxide (125 mg, 1.3 mmol) was added to a stirred solution of 4-fluoro-1-bromobenzene (176 mg, 1.0 mmol), 1-(4-fluorophenyl)-2-methoxyethanone (Method 23; 185 mg), 2-dicyclohexylphosphino-2'-(*N,N*-dimethylamino)-biphenyl (8 mg, 0.02 mmol) and palladium acetate (2.2 mg, 0.01 mmol) in dry toluene (1 ml) under argon. The resultant mixture was heated at 80°C for 16 hours, cooled to room temperature and partitioned between water (10.0 ml) and ether (25 ml). The aqueous layer was extracted with ether (2x10 ml), the combined ether extracts washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in hexane as eluent to give the title compound as a solid (178 mg, 0.68 mmol). NMR: 3.4 (s, 3H), 5.4 (s, 1H), 7.0 (m, 4H), 7.4 (dd, 2H) and 8.0 (dd, 2H).

**Examples 38-41**

The procedure described in Example 37 was repeated using the appropriate bromobenzene to replace the 4-fluoro-1-bromobenzene to obtain the compounds described below.

100804-2

- 72 -

Ex	Compound	NMR
38	( $\alpha$ -Methoxy-4-methylbenzyl)-(4-fluorophenyl)-ketone	2.3 (s, 3H), 3.4 (s, 3H), 5.4 (s, 1H), 7.0 (dd, 2H), 7.1 (d, 2H), 7.3 (d, 2H), 8.0 (dd, 2H)
39	( $\alpha$ -Methoxy-4-methoxybenzyl)-(4-fluorophenyl)-ketone	3.4 (s, 3H), 3.8 (s, 3H), 5.4 (s, 1H), 6.8 (d, 2H), 7.0 (dd, 2H), 7.3 (d, 2H), 8.0 (dd, 2H)
40	( $\alpha$ -Methoxy-4-[N,N-dimethylsulphamoyl]benzyl)-(4-fluorophenyl)-ketone	2.7 (s, 6H), 3.5 (s, 3H), 5.4 (s, 1H), 7.1 (dd, 2H), 7.6 (d, 2H), 7.8 (d, 2H), 8.0 (dd, 2H)
41	[ $\alpha$ -Methoxy-4-(methoxymethyl)benzyl]-(4-fluorophenyl)-ketone	3.2 (s, 2H), 3.4 (s, 3H), 3.45 (s, 3H), 5.4 (s, 1H), 7.0 (m, 4H), 7.4 (dd, 2H), 8.0 (dd, 2H)

**Reference Example 21****(4-Methyl- $\alpha$ -hydroxybenzyl)-(4-chlorophenyl)-ketone**

5 A solution of sodium methoxide in methanol (10.0ml of a 0.5 mol solution, 5.0 mmol) was added to a stirred solution of 2-bromo-1-(4-chlorophenyl)-2-(4-methylphenyl)-ethan-1-one (323 mg, 1.0 mmol) in methanol (10 ml) during 30 minutes at ambient temperature. The resultant mixture was stirred at ambient temperature for 3 hours and then quenched with 1M hydrochloric acid (5.0 ml). The methanol was evaporated and the aqueous residue treated with  
 10 ether (20 ml). The ether layer was separated, washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in hexane as eluent to give the title compound as a solid (215 mg, 0.78 mmol). NMR: 2.3 (s, 3H), 4.4 (d, 1H), 5.8 (d, 1H), 7.1-7.2 (m, 4H), 7.4 (dd, 2H) and 7.8 (dd, 2H).

**15 Example 42****(4-Fluorophenyl)-[ $\alpha$ -(5-chloropyrimidin-2-yloxy)-4-fluorobenzyl]-ketone**

To a stirred solution of 5-chloro-2-hydroxypyrimidine (130 mg, 1.0 mmol), 4,4'-difluorobenzoin (372 mg, 1.5 mmol) and triphenylphosphine (524 mg, 2 mmol) in dry THF (10 ml) was added a solution of di-isopropylazodicarboxylate (445 mg, 2.2 mmol) at 0°C  
 20 under argon. The resultant mixture was stirred at ambient temperature for 16 hours,

100804-2

- 73 -

partitioned between water (25 ml) and ether (25 ml). The aqueous layer was extracted with ether (25 ml), and the combined ether extracts were washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 50% EtOAc in hexane as eluent to give the title compound as a solid (74 mg, 0.21 mmol). NMR: 7.1 (m, 4H), 7.3 (m, 2H), 7.4 (d, 1H), 7.5 (s, 1H), 8.0 (m, 2H) and 8.5 (d, 1H);  $m/z$  359 (M-H).

### Example 43

#### ( $\alpha$ -Hydroxy-4-methoxybenzyl)-(naphth-2-yl)-ketone

2-Naphthaldehyde (3.75g, 24mmol) was dissolved in DCM (50ml) and zinc diiodide (250mg) was added. This was stirred at room temperature under argon and trimethylsilyl cyanide (6.65ml, 25mmol) was added via syringe. The reaction mixture was stirred overnight. The solvent was removed *in vacuo* to leave an orange oil. LDA was prepared by adding diisopropylamine (3.35ml, 24mol) in THF (25ml) and cooling to  $-60^{\circ}\text{C}$ , before adding *n*-butyl lithium (1.54ml) under argon. This was stirred for 15 mins before adding the orange oil – the cyanohydrin – in THF (20ml) and stirring at  $-60^{\circ}\text{C}$  for 30 mins. Para-anisaldehyde (2.92ml, 24 mmol) in THF (15ml) was added and the reaction was allowed to stir and warm up to room temperature overnight. Saturated aqueous ammonium chloride (65ml) was added to the reaction mixture followed by ether (100ml). The organic phase was separated, washed with saturated ammonium chloride, dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to give an orange oil. This was taken up in methanol (30ml) and 1M sulphuric acid (10ml) was added. The reaction mixture was left to stand overnight. The pH was adjusted to pH 7 - 8, and the mixture was concentrated and extracted with DCM. The organic layers were combined, washed, dried and evaporated to give an orange oil, which was purified by column chromatography (EtOAc:hexane, 10:1) to give a pale yellow solid (56mg, 0.8%). Mp  $121 - 128^{\circ}\text{C}$ ; NMR (200MHz,  $\text{DMSO}-d_6$ ): 3.65 (s, 3H), 5.8 – 5.9 (bs, 1H), 6.20 (s, 1H), 6.80 and 7.35 (AB q, 4H), 7.60 – 7.90 (m, 4H).

### Reference Example 22

#### ( $\alpha$ -Hydroxy-4-methoxybenzyl)-(4-methoxyphenyl)-ketone

Anisaldehyde (20g) was dissolved in methanol (25ml) and water (16ml). Potassium cyanide (4g) was added and the mixture was refluxed for 2 hours. Further potassium cyanide (4g) was added and the reaction refluxed for a further 2 hours. On allowing to stand, an oil separated. The solvent was removed *in vacuo* and the residue was taken up in water, and



100804-2

- 74 -

extracted with ether. The extracts were combined, washed with water, dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to give an oil. This was extracted with hot petroleum ether 60 – 80°C to remove the anisaldehyde, and the residue triturated with ethanol to give a solid (1.6g). This was recrystallized from aqueous ethanol to give the product (820mg). Mp 110 – 112°C;  
 5 m/z 272 ( $\text{M}^+$ ).

#### Examples 44-45 and Reference Example 23

Following the procedure of Reference Example 22 using the appropriate starting materials the following compounds were prepared.

Ex	Compound	Data
RE 23	( $\alpha$ -Hydroxybenzyl)-(4-methoxyphenyl)-ketone	Mp 105.5 – 106°C (lit. 106°C); m/z 242 ( $\text{M}^+$ )
44	(1-Naph-2-yl-1-hydroxymethyl)-[4-( <i>N,N</i> -dimethylamino)phenyl]-ketone	Mp 128 - 132°C; NMR (200MHz, $\text{DMSO-d}_6$ ): 2.95 (s, 6H), 5.85 (bs, 1H), 6.10 (s, 1H), 6.60 (AB q, 2H), 7.50 (m, 3H), 7.90 (m, 6H)
45 <sup>1</sup>	( $\alpha$ -Hydroxy-3,4-dichlorobenzyl)-(3,4-dichlorophenyl)-ketone	Mp 100 - 102°C; m/z 332 ( $\text{M}^+$ )

10 <sup>1</sup> This compound was prepared with sodium cyanide, not potassium cyanide.

#### Example 46

##### [ $\alpha$ -Hydroxy- $\alpha$ -(*N,N*-diisopropylaminomethyl)benzyl]-(phenyl)-ketone

Diisopropylamine (11.6g, 115mmol) was added to a solution of 2-hydroxy-1,2-  
 15 diphenyl-ethanone (21g, 100mmol) and 40% aqueous formaldehyde (10ml, 140mmol) in ethanol (40ml) and the mixture was heated under reflux for 2 hours. The mixture was cooled to room temperature and partitioned between water (200ml) and ether (600ml). The ether layer was washed with water (2x200ml) and extracted with 1M hydrochloric acid (3x150ml). The combined acidic extracts were basified with concentrated aqueous sodium hydroxide  
 20 solution and extracted with ether (3x150ml). The combined ether extracts were dried, treated with hydrogen chloride in ethanol until acidic and evaporated to dryness. The residue was crystallised from ethanol to give the title compound as a solid (3.9g, 10.8mmol). M/z 325 ( $\text{M}^+$ ).

25

100804-2

- 75 -

**Example 47****(2-Thien-2-ylethyl)-(4-chlorophenyl)-ketone**

A solution of 4-chlorophenyl magnesium bromide in ether (6.0ml of a 1.0 mol solution, 6.0 mmol) was added to a stirred solution of *N*-methoxy-*N*-methyl-2-thienylethanamide (Method 2; 398 mg, 2.0 mmol) in THF (20 ml) at 0°C. The resultant mixture was stirred at ambient temperature overnight and then quenched with ethanol (50 ml). The resultant mixture was evaporated to dryness and the residue partitioned between water (50 ml) and ether (100 ml). The ether layer was separated, washed with brine, dried and evaporated to dryness. The residue was purified by column chromatography using 5% EtOAc in hexane as eluent to give the title compound as a solid (250 mg, 1.0 mmol). NMR: 3.3 (m, 4H), 6.8 (dd, 1H), 6.9 (dd, 1H), 7.1 (dd, 1H), 7.4 (d, 2H), 7.9 (d, 2H).

**Reference Examples 24-26 and Examples 48-50**

The procedure described in Example 47 was repeated using the appropriate *N*-methoxy-*N*-methyl amide to replace the *N*-methoxy-*N*-methyl-2-thienylethanamide and the appropriate Grignard or lithium reagent to replace the 4-chlorophenyl magnesium bromide to obtain the compounds described below.

Ex	Compound	NMR
RE 24	(4-Fluorophenethyl)-(4-fluorophenyl)-ketone	3.0 (t, 2H), 3.2 (t, 2H), 6.9 (dd, 2H), 7.1 (dd, 2H), 7.2 (dd, 2H), 8.0 (dd, 2H)
RE 25	(4-Chlorophenethyl)-(4-fluorophenyl)-ketone	3.0 (t, 2H), 3.2 (t, 2H), 7.0 (dd, 2H), 7.1 (dd, 2H), 7.2 (dd, 2H), 8.0 (dd, 2H)
RE 26	(2-Thien-2-ylethyl)-(4-fluorophenyl)-ketone	3.3 (m, 4H), 6.8 (d, 1H), 6.9 (dd, 1H), 7.1 (m, 3H), 8.0 (dd, 2H)
48	(2-Thien-2-ylethyl)-(4-methylphenyl)-ketone	2.4 (s, 3H), 3.3 (m, 4H), 6.8 (d, 1H), 6.9 (dd, 1H), 7.1 (dd, 1H), 7.2 (dd, 2H), 7.8 (dd, 2H)
49	(4-Chlorophenethyl)-(thiazol-2-yl)-ketone	3.0 (t, 2H), 3.5 (t, 2H), 7.2 (m, 4H), 7.6 (d, 1H), 8.0 (d, 1H)
50	(2-Thien-2-ylethyl)-(thiazol-2-yl)-ketone	3.3 (t, 2H), 3.6 (t, 2H), 6.9 (dd, 2H), 7.1 (d, 1H), 7.6 (d, 1H), 8.0 (d, 1H)

100804-2

- 76 -

**Reference Examples 27-28 and Examples 51-53**

The following compounds were made by the procedure of J.Med.Chem.; EN; 30; 12; 1987; 2232-2239.

Ex	Compound	M/z	NMR
RE 27	(Morpholinosulphonylmethyl)-(4-fluorophenyl)-ketone	286	3.3 (dd, 4H), 3.7 (dd, 4H), 4.5 (s, 2H), 7.2 (m, 2H), 8.0 (m, 2H)
51	(Piperidin-1-ylsulphonylmethyl)-(4-fluorophenyl)-ketone	284	1.6 (m, 6H), 3.3 (m, 4H), 4.5 (s, 2H), 7.2 (m, 2H), 8.0 (m, 2H)
52	[4-(4-Fluorophenyl)piperidin-1-ylsulphonylmethyl]-(4-fluorophenyl)-ketone	378	
RE 28	[N-Methylanilinosulphonylmethyl]-(phenyl)-ketone	288 (M-H) <sup>+</sup>	3.35 (s, 3H), 4.60 (s, 2H), 7.35 (m, 1H), 7.40 (m, 2H), 7.50 (m, 4H), 7.60 (m, 1H), 8.00 (d, 2H)
53	(Morpholinosulphonylmethyl)-(phenyl)-ketone	284 (M-H) <sup>+</sup>	2.70 (m, 4H), 3.65 (m, 2H), 4.60 (s, 2H), 7.55 (t, 2H), 7.70 (t, 1H), 8.05 (d, 2H)

**5 Example 54****(4-Fluorophenyl)-[N-(cyclohexyl)-N-(isopropyl)sulphamoylmethyl]-ketone**

To a stirred solution of *N*-(isopropyl)-*N*-(mesyl)cyclohexylamino (Method 12; 225mg, 1.03mmol) in anhydrous THF (5ml) at -20°C was added a 1M solution of lithium bis(trimethylsilyl)amide (2.06ml, 2.06mmol). The reaction was stirred at -20°C for 30 mins  
 10 before the addition of a solution of methyl-4-fluoro benzoate (206mg, 1.33mmol) in anhydrous THF (2ml). The reaction was allowed to warm to room temperature and then stirred at this temperature for 1 hour. The reaction was quenched with saturated ammonium chloride (~5ml) and the organic layer was separated. The aqueous layer was reextracted with EtOAc. The combined organic layers were washed with brine then dried (MgSO<sub>4</sub>), filtered  
 15 and evaporated to yield an oil. This oil was purified by column chromatography (DCM to 5%MeOH/DCM) to yield the product as an oil which crystallised on standing (153mg, 44%).

100804-2

- 77 -

NMR: 1.00 (m, 1H), 1.20 (m, 3H), 1.25 (d, 6H), 1.55 (m, 1H), 1.65 (m, 2H), 1.75 (m, 3H), 3.15 (m, 1H), 3.65 (m, 1H), 4.45 (s, 2H), 7.10 (t, 2H), 8.05 (m, 2H); m/z: 340 (M-H)<sup>+</sup>.

### Examples 55-59 and Reference Example 29

- 5 The procedure described in Example 54 was repeated using the appropriate starting materials to obtain the compounds described below.

Ex	Compound	NMR	M/z	SM
55	(4-Fluorophenyl)-[N-(4-chlorophenyl)-N-(methyl)sulphamoylmethyl]-ketone	3.30 (s, 3H), 4.50 (s, 2H), 7.15 (t, 2H), 7.35 (d, 2H), 7.50 (d, 2H), 8.05 (m, 2H)	340 (M-H) <sup>+</sup>	Method 15
56	(4-Fluorophenyl)-[N-(pyrid-2-yl)-N-(methyl)sulphamoylmethyl]-ketone	3.45 (s, 3H), 5.05 (s, 2H), 7.15 (t, 3H), 7.25 (d, 1H), 7.75 (t, 1H), 8.05 (m, 2H), 8.40 (m, 1H)	307 (M-H) <sup>+</sup>	Method 13
57	(4-Fluorophenyl)-[N-(4-methoxyphenyl)-N-(ethyl)sulphamoylmethyl]-ketone	1.05 (t, 3H), 3.65 (q, 2H), 3.85 (s, 3H), 4.55 (s, 2H), 6.95 (d, 2H), 7.20 (t, 2H), 7.45 (d, 2H), 8.10 (m, 2H)	350 (M-H) <sup>+</sup>	Method 17
58 <sup>1</sup>	(4-Fluorophenyl)-[N-(4-chlorophenyl)sulphamoylmethyl]-ketone	4.50 (s, 2H), 7.00 (br s, 1H), 7.20 (t, 2H), 7.30 (m, 4H), 7.95 (m, 2H)	326 (M-H) <sup>+</sup>	Method 18
59	(4-Fluorophenyl)-[N-(4-methoxyphenyl)-N-(methyl)sulphamoylmethyl]-ketone	3.30 (s, 3H), 3.80 (s, 3H), 4.55 (s, 2H), 6.90 (d, 2H), 7.15 (t, 2H), 7.45 (d, 2H), 8.05 (m, 2H)	336 (M-H) <sup>+</sup>	Method 16
RE 29	(4-Fluorophenyl)-(4-methylpiperazin-1-yl)sulphamoylmethyl]-ketone	2.30 (s, 3H), 2.45 (m, 4H), 3.35 (m, 4H), 4.50 (s, 2H), 7.20 (t, 2H), 8.10 (m, 2H)	299 (M-H) <sup>+</sup>	Method 14

<sup>1</sup> In this example 3 equivalents of lithium bis(trimethylsilyl)amide were used and the final product was crystallised from ether

100804-2

- 78 -

**Example 60****(4-Bromophenylsulphonylmethyl)-(4-cyanophenyl)-ketone**

To a stirred solution of methyl 4-cyanobenzoate (150mg, 0.93mmol) and 4-bromophenyl methyl sulphone (200mg, 0.84mmol) in 1,2-dimethoxyethane (10ml) was added sodium hydride (40%) (120mg, 3mmol). The reaction was warmed to 85°C and stirred at this temperature for 6 hours. The reaction was allowed to cool to room temperature and then quenched with water (~50ml). The solution was transferred to a separating funnel and washed with ether, the layers were separated and the organic layer was extracted with 1M sodium hydroxide solution. The aqueous layers were combined and acidified to ~pH3 with concentrated hydrochloric acid. The resulting suspension was extracted with DCM (2x50ml), the organic layers were combined and washed with brine then dried (MgSO<sub>4</sub>), filtered and evaporated to yield an oil. The oil was purified by column chromatography (10g silica, DCM) to yield a clear oil which crystallised on standing. NMR: 4.65 (s, 2H), 7.65 (m, 4H), 7.75 (d, 2H), 8.00 (d, 2H); m/z 363 (M-H)<sup>+</sup>.

15

**Examples 61-71 and Reference Example 30**

The procedure described in Example 19 was repeated using the appropriate starting materials.

Ex	Compound	M/z	NMR
61	(4-Bromophenylsulphonylmethyl)-(4-trifluoromethylphenyl)-ketone	406 (M-H) <sup>+</sup>	4.75 (s, 2H), 7.70 (m, 4H), 7.80 (d, 2H), 8.10 (d, 2H)
62	(4-Fluorophenylsulphonylmethyl)-(4-trifluoromethylphenyl)-ketone	345 (M-H) <sup>+</sup>	4.70 (s, 2H), 7.20 (m, 2H), 7.70 (d, 2H), 7.85 (m, 2H), 8.00 (d, 2H)
63	(Thien-2-ylsulphonylmethyl)-(thien-2-yl)-ketone	271 (M-H) <sup>+</sup>	4.70 (s, 2H), 7.15 (m, 2H), 7.75 (br m, 4H)
64	(Thien-2-ylsulphonylmethyl)-(4-cyanophenyl)-ketone	290 (M-H) <sup>+</sup>	4.85 (s, 2H), 7.15 (m, 1H), 7.65 (m, 1H), 7.80 (m, 3H), 8.10 (d, 2H)
65	(Thien-2-ylsulphonylmethyl)-(4-trifluoromethylphenyl)-ketone	333 (M-H) <sup>+</sup>	4.80 (s, 2H), 7.10 (m, 1H), 7.60 (d, 1H), 7.70 (m, 3H), 8.00 (d, 2H)
66	(4-Bromophenylsulphonylmethyl)-(thien-2-yl)-ketone	344 (M-H) <sup>+</sup>	4.60 (s, 2H), 7.20 (m, 1H), 7.75 (br m, 6H)
67	(4-Methylphenylsulphonylmethyl)-(4-cyanophenyl)-ketone	298 (M-H) <sup>+</sup>	4.75 (s, 2H), 7.35 (d, 2H), 7.75 (m, 4H), 8.05 (d, 2H)

100804-2

- 79 -

68	(4-Fluorophenylsulphonylmethyl)-(4-fluorophenyl)-ketone	295 (M-H) <sup>-</sup>	4.65 (s, 2H), 7.20 (m, 4H), 7.90 (m, 2H), 8.00 (m, 2H)
69	(Thien-2-ylsulphonylmethyl)-(4-fluorophenyl)-ketone	283 (M-H) <sup>-</sup>	4.80 (s, 2H), 7.20 (m, 3H), 7.70 (m, 2H), 8.00 (m, 2H)
70	(Thien-2-ylsulphonylmethyl)-(fur-2-yl)-ketone	255 (M-H) <sup>-</sup>	4.70 (s, 2H), 6.60 (m, 1H), 7.15 (m, 1H), 7.35 (m, 1H), 7.60 (s, 1H), 7.70 (d, 1H), 7.75 (d, 1H)
RE 30	(4-Methylphenylsulphonylmethyl)-(fur-2-yl)-ketone	263 (M-H) <sup>-</sup>	2.45 (s, 3H), 4.55 (s, 2H), 6.60 (m, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 7.80 (d, 2H)
71 <sup>1</sup>	(4-Methoxyphenylsulphonylmethyl)-(fur-2-yl)-ketone	279 (M-H) <sup>-</sup>	3.90 (s, 3H), 4.55 (s, 2H), 6.60 (m, 1H), 7.00 (d, 2H), 7.30 (m, 1H), 7.60 (s, 1H), 7.80 (d, 2H)

<sup>1</sup> In this example the sulphone used was 4-fluorophenyl methyl sulphone, the fluorine is displaced by methoxide during the reaction.

#### Example 72

##### 5 (Phenylsulphonylmethyl)-(pyrid-2-yl)-ketone

A solution of methylphenyl sulphone (3g, 19.2mmol) in THF was added dropwise to a solution of lithium diisopropyl amine (2.7ml diisopropylamine and 12ml of 1.6M *n*-butyl lithium) in THF under argon at -78°C. The resultant pink-orange solution was stirred at -78°C for 15 mins. A solution of pyridine 2-methylcarboxylate (1.32g, 9.6mmol) in THF was added.

- 10 The reaction mixture was stirred at -78°C for 2 hours, then allowed to warm to room temperature and was stirred overnight. The reaction was quenched with water, filtered and evaporated to dryness to give an oil. This oil was taken up in EtOAc and purified with flash chromatography (2:1 EtOAc:petrol) to give the product which was recrystallized from EtOAc/petrol to give the solid (840mg). Mp 101 - 103°C; m/z 261 (M<sup>+</sup>).

15

#### Reference Example 31

##### (4-Bromophenyl)-(4-methylphenylsulphonylmethyl)-ketone

- 4-Sulphotoluene sodium salt (21.4g) and 2,4'-dibromoacetophenone (27.8g) were mixed in ethanol (100ml) and refluxed overnight. The reaction was cooled after which the product crystallised. The crystals were filtered and dried. The filtrate was evaporated to give
- 20

100804-2

- 80 -

another crop of crystals. The first crop were triturated with water (200ml) and filtered and combined with the second crop. The combined mass was taken up in ethanol (200ml) and refluxed until all material dissolved. The resulting crystals were separated by filtration and dried. NMR (400MHz, DMSO- $d_6$ ): 2.30 (s, 3H), 5.15 (s, 2H), 7.30 (d, 2H), 7.60 (m, 4H), 7.75 (d, 2H);  $m/z$  354.

#### Examples 73-76 and Reference Examples 32-34

The following compounds were made by the procedure of Syn.Lett.; EN; 10; 2000; 1500 - 1502 (except 1.2eq of NaI was added to the reaction mixture) using the appropriate 10 starting materials.

Ex	Compound	M/z	NMR (DMSO- $d_6$ )
73	[ $\alpha$ -(2-Methylthiazol-4-ylmethyl)benzyl]-(phenyl)-ketone	308	2.50 (s, 3H), 3.05 (dd, 1H), 3.50 (dd, 1H), 5.35 (t, 1H), 6.95 (s, 1H), 7.15 (m, 1H), 7.25 (m, 2H), 7.35 (m, 2H), 7.45 (t, 2H), 7.55 (t, 1H), 8.00 (d, 2H)
74	[ $\alpha$ -(2-Chlorothiazol-5-ylmethyl)benzyl]-(phenyl)-ketone	328	3.25 (dd, 1H), 3.55 (dd, 1H), 5.15 (t, 1H), 7.15 (m, 1H), 7.30 (m, 5H), 7.45 (t, 2H), 7.55 (t, 1H), 8.00 (d, 2H)
RE 32	[ $\alpha$ -(Cyanomethyl)benzyl]-(phenyl)-ketone	493 [2M+Na]	(CDCl <sub>3</sub> ): 2.90 (m, 1H), 3.10 (m, 1H), 4.85 (t, 1H), 7.30 (br m, 7H), 7.50 (t, 1H), 7.90 (d, 2H)
RE 33	[ $\alpha$ -(Benzyl)benzyl]-(phenyl)-ketone	287	3.00 (m, 1H), 3.45 (m, 1H), 5.20 (t, 1H), 7.10 (m, 3H), 7.15 (m, 3H), 7.30 (br m, 4H), 7.40 (t, 2H), 7.50 (t, 1H), 8.00 (d, 2H)
RE 34	[ $\alpha$ -(Propyl)benzyl]-(phenyl)-ketone	239	0.90 (t, 3H), 1.25 (m, 2H), 1.80 (m, 1H), 2.15 (m, 1H), 4.55 (t, 1H), 7.20 (m, 1H), 7.30 (m, 4H), 7.40 (t, 2H), 7.45 (m, 1H), 7.95 (d, 2H)
75 1	(5-Methylfur-2-yl)-[2-(4-chlorophenyl)-1-(pyrazin-2-yl)ethyl]-ketone	327	

100804-2

- 81 -

76	(4-Fluorophenyl)-[2-(2-chlorothiazol5-yl)-1-(thien-3-yl)ethyl]-ketone	352	3.25 (dd, 1H), 3.60 (dd, 1H), 4.85 (t, 1H), 6.95 (d, 1H), 7.15 (t, 3H), 7.20 (s, 1H), 7.30 (m, 1H), 7.95 (m, 2H)
----	---	-----	--

<sup>1</sup> This example required purification by prep LCMS after column chromatography.

### Reference Example 35

#### 5 (N-Methyl-4-methylanilinosulphonylmethyl)-(4-chlorophenyl)-ketone

To a stirred solution of *N*-methyl-4-methylanilinosulphonylmethyl (EP 495594; 199mg, 1.0mmol) in dry THF (1 ml) at -78°C was added a solution of 1.6M *n*-butyl lithium in hexane (1.25 ml, 2.0mmol). The reaction was stirred at room temperature for 1 hour then cooled to -78°C and treated with a solution of methyl-4-chlorobenzoate (170mg, 1.0mmol) in  
 10 dry THF (1 ml). The mixture was stirred at -78°C for 2 hours then stirred at room temperature for 1 hour. The reaction was quenched with saturated ammonium chloride solution (5 ml) and water (5 ml) and extracted with ether (2x20 ml). The combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and evaporated to dryness. The residue was purified by column chromatography using 10% EtOAc in hexane as eluent to give a solid which was  
 15 crystallised from ether/hexane to give the title compound (280 mg, 0.83mmol). NMR: 2.4 (s, 3H), 3.3 (s, 3H), 4.5 (s, 2H), 7.2 (m, 2H), 7.4 (m, 4H), 8.00 (d, 2H).

### Examples 77-78

The procedure described in Reference Example 35 was repeated using the appropriate  
 20 sulphonamide and ester to yield the desired product.

Ex	Compound	NMR / m/z
77 <sup>1</sup>	( <i>N</i> -Methyl-4-methoxyanilinosulphonylmethyl)-(4-chlorophenyl)-ketone	NMR: 3.3 (s, 3H), 3.8 (s, 3H), 4.5 (s, 2H), 6.9 (m, 2H), 7.4 (m, 4H), 8.00 (d, 2H)
78 <sup>2,3</sup>	(4-Bromo-2-methoxycarbonylanilinosulphonylmethyl)-(4-bromo-2-mesylaminophenyl)-ketone	582

<sup>1</sup> *N*-methyl-4-methoxyanilinosulphonylmethyl: Advanced Synthesis and Catalysis 2001, 343 (1), 71-74; synthesized by reaction of methanesulphonyl chloride and 4-methoxy-*N*-methylaniline in pyridine

<sup>2</sup> The base used was LDA.



100804-2

- 82 -

<sup>3</sup> Condensation reaction with 4-bromo-2-methoxycarbonylanilinosulphonylmethyl.

### Example 79

#### [ $\alpha$ -(4-Methoxycarbonylbenzyl)benzyl]-(phenyl)-ketone

5 To deoxybenzoin (50 mg, 0.25 mmol) in THF (2ml) at 0°C under a nitrogen atmosphere was added dropwise a 1M solution of lithium bis(trimethylsilyl)amide in THF (0.28 ml, 0.28 mmol). The reaction was stirred at 0°C for 3 hours 30 mins before being added dropwise to a solution of methyl 4-(bromomethyl)benzoate (229 mg, 0.28 mmol) in THF (2 ml) at 0°C under a nitrogen atmosphere. The reaction was stirred in the melting ice bath for 16  
10 hours. Water (5 ml) was added slowly to the reaction, which was then extracted with DCM (3x15 ml). The combined organic layers were concentrated *in vacuo*. The crude product was chromatographed on Kieselgel 60, eluting with 15% EtOAc in *iso*-hexane, to give the product as a white solid (57 mg, 66%). NMR (300MHz, DMSO- $d_6$ ) 3.05 (1H, dd), 3.45 (1H, dd), 3.80 (3H, s), 5.25 (1H, t), 7.35 (10H, m), 7.75 (2H, d), 7.95 (2H, d); *m/z* 345.

15

#### Reference Examples 36-37 and Examples 80-91

The procedure described in Example 79 was repeated using the appropriate starting materials.

Ex	Compound	M/z	NMR (300MHz, DMSO- $d_6$ )
RE 36 <sup>A</sup>	( $\alpha$ -Methylbenzyl)-(4-chlorophenyl)-ketone	245	1.40 (3H, d), 4.90 (1H, q), 7.30 (5H, m), 7.50 (2H, d), 7.95 (2H, d)
80	[ $\alpha$ -(Benzyl)benzyl]-(5-bromothiien-2-yl)-ketone	371	3.00 (1H, dd), 3.40 (1H, dd), 5.05 (1H, t), 7.25 (11H, m), 7.95 (1H, d)
RE 37	[ $\alpha$ -(Benzyl)benzyl]-(4-chlorophenyl)-ketone	321	3.00 (1H, dd), 3.45 (1H, dd), 5.20 (1H, t), 7.20 (10H, m), 7.50 (2H, d), 8.00 (2H, d)
81	(1-Phenyl-3-morpholinoprop-2-yl)-(thien-2-yl)-ketone	316	2.30 (2H, m), 2.45 (3H, m), 2.65 (1H, m), 2.85 (2H, m), 3.40 (4H, m), 3.95 (1H, m), 7.15 (6H, m), 7.90 (2H, m)

100804-2

- 83 -

82	[ $\alpha$ -(Benzyl)benzyl]-(thien-2-yl)-ketone	293	3.00 (1H, dd), 3.40 (1H, dd), 5.05 (1H, t), 7.15 (7H, m), 7.25 (2H, t), 7.40 (2H, d), 7.90 (1H, d), 8.05 (1H, d)
83	[ $\alpha$ -(Pyrid-3-ylmethyl)benzyl]-(phenyl)-ketone	288	3.05 (1H, dd), 3.40 (1H, dd), 5.25 (1H, t), 7.35 (10H, m), 8.00 (2H, d), 8.35 (2H, m)
84	[ $\alpha$ -(Pyrid-2-ylmethyl)benzyl]-(phenyl)-ketone	288	3.15 (1H, dd), 3.65 (1H, dd), 5.50 (1H, dd), 7.35 (11H, m), 8.00 (2H, d), 8.35 (1H, d)
85	[ $\alpha$ -(3-Methoxycarbonylbenzyl)benzyl]-(phenyl)-ketone	313 [M-OMe] <sup>+</sup>	3.05 (1H, dd), 3.45 (1H, dd), 3.80 (3H, s), 5.25 (1H, t), 7.35 (10H, m), 7.70 (1H, d), 7.80 (1H, s), 7.95 (2H, d)
86	[ $\alpha$ -(Pyrid-4-ylmethyl)benzyl]-(phenyl)-ketone	288	3.05 (1H, dd), 3.40 (1H, dd), 5.30 (1H, t), 7.35 (10H, m), 8.00 (2H, d), 8.35 (2H, d)
87	[ $\alpha$ -(2-Ethoxycarbonylbenzyl)benzyl]-(phenyl)-ketone	345	1.25 (3H, t), 3.05 (1H, dd), 3.45 (1H, dd), 4.25 (2H, q), 5.25 (1H, t), 7.35 (10H, m), 7.75 (2H, d), 7.95 (2H, d)
88	[ $\alpha$ -(2-Nitrobenzyl)benzyl]-(phenyl)-ketone	332	3.25 (1H, dd), 3.65 (1H, dd), 5.20 (1H, t), 7.35 (11H, m), 7.85 (1H, d), 7.95 (2H, d)
89	[ $\alpha$ -(3-Nitrobenzyl)benzyl]-(phenyl)-ketone	332	3.15 (1H, dd), 3.35 (1H, dd), 5.30 (1H, t), 7.40 (10H, m), 8.00 (4H, m)
90	[ $\alpha$ -(3-Nitro-6-methoxybenzyl)benzyl]-(phenyl)-ketone	362	3.10 (1H, dd), 3.40 (1H, dd), 3.85 (3H, s), 5.15 (1H, t), 7.15 (6H, m), 7.45 (2H, m), 7.95 (5H, m)

100804-2

- 84 -

90	[ $\alpha$ -(5-Nitrofur-2-ylmethyl)benzyl]- (phenyl)-ketone	3.20 (1H, dd), 3.35 (1H, dd), 5.35 (1H, t), 6.50 (1H, d), 7.35 (9H, m), 8.00 (2H, d)
----	--	--

<sup>A</sup> Methyl iodide was the alkylating reagent.**Example 92****5 (4-Cyanophenoxymethyl)-(4-chlorophenyl)-ketone**

2-Bromo-4'-chloroacetophenone (500mg, 2.15mmol), 4-cyanophenol (256.4mg, 2.15mmol) and potassium carbonate (297.4mg, 2.15mmol) were placed in acetone and the reaction mixture was stirred and heated at reflux overnight. On cooling, the solvent was evaporated *in vacuo* and the residue was partitioned between EtOAc and water. The organic  
10 layer was separated, dried (MgSO<sub>4</sub>) and the organics removed *in vacuo* to give a brown solid. This was triturated with a 1:1 mixture of EtOAc and hexane to give a white solid, which was collection by filtration, 327.7mg, 56%. NMR (300MHz): 5.25 (s, 2H), 6.95 (d, 2H), 7.45 (d, 2H), 7.55 (d, 2H), 7.90 (d, 2H); m/z 270 for (M-H)<sup>+</sup>.

15

100804-2

- 85 -

**Examples 93-125 and Reference Examples 38-42**

The procedure described in Example 92 was repeated using the appropriate starting materials.

Ex	Compound	M/z	NMR
93	(4-Ethoxyphenoxyethyl)-(4-chlorophenyl)-ketone	291	1.30 (t, 3H), 3.90 (q, 2H), 5.05 (s, 2H), 6.75 (m, 4H), 7.40 (d, 2H), 7.85 (d, 2H)
94	(4-Phenylphenoxyethyl)-(4-chlorophenyl)-ketone	323	5.05 (s, 2H), 6.90 (d, 2H), 7.25 (t, 1H), 7.35 (t, 2H), 7.45 (m, 6H), 7.90 (d, 2H)
95	(4-Mesylphenoxyethyl)-(4-chlorophenyl)-ketone	323 (M-H) <sup>+</sup>	2.95 (s, 3H), 5.25 (s, 2H), 6.95 (d, 2H), 7.45 (d, 2H), 7.80 (d, 2H), 7.85 (d, 2H)
96	(4-Fluoro-3-chlorophenoxyethyl)-(4-chlorophenyl)-ketone	297 (M-H) <sup>+</sup>	5.20 (s, 2H), 6.80 (m, 1H), 6.95 - 7.05 (m, 2H), 7.50 (d, 2H), 7.90 (d, 2H)
97	(4-Fluoro-2-chlorophenoxyethyl)-(4-chlorophenyl)-ketone	297 (M-H) <sup>+</sup>	5.20 (s, 2H), 6.80 (m, 2H), 7.10 (m, 1H), 7.45 (dd, 2H), 7.95 (dd, 2H)
98	(4-Cyanomethylphenoxyethyl)-(4-chlorophenyl)-ketone	286	3.70 (s, 2H), 5.20 (s, 2H), 6.90 (d, 2H), 7.20 (d, 2H), 7.50 (d, 2H), 7.90 (d, 2H)
99	[4-(2-Thiazolin-2-yl)phenoxyethyl]-(4-chlorophenyl)-ketone	332	3.40 (t, 2H), 4.40 (t, 2H), 5.25 (s, 2H), 6.90 (d, 2H), 7.50 (d, 2H), 7.75 (d, 2H), 7.95 (d, 2H)
100	(4-Cyanophenoxyethyl)-(2,4-dichlorophenyl)-ketone	305	5.50 (s, 2H), 7.15 (dt, 2H), 7.65 (dd, 1H), 7.75 - 7.85 (m, 3H), 7.95 (d, 1H)
101	(2-Methylpyrid-5-yloxyethyl)-(phenyl)-ketone	228	2.38 (s, 3H), 5.62 (s, 2H), 7.14 (d, 1H), 7.30 (m, 3H), 7.65 (d, 1H), 7.68 (d, 1H), 8.0 (d, 1H), 8.19 (d, 1H)

100804-2

- 86 -

102	(2-Carbamoylphenoxyethyl)-(4-bromophenyl)-ketone	334	7.20 (td, 1H), 7.40 (d, 1H), 7.60 (td, 1H), 7.80 (bs, 1H), 7.95 (dt, 2H), 8.05 – 8.15 (m, 3H), 8.45 (bs, 1H)
103 1	(4-fluorophenoxyethyl)-(4-chlorophenyl)-ketone	264	5.40 (s, 2H), 6.80 – 6.90 (m, 2H), 6.95 – 7.05 (m, 2H), 7.55 (m, 2H), 7.90 (m, 2H)
RE 38	(Naphth-2-yloxyethyl)-(phenyl)-ketone	261	5.65 (s, 2H), 7.25 (dd, 1H), 7.30 – 7.35 (m, 2H), 7.45 (td, 1H), 7.60 (t, 2H), 7.65 – 7.75 (m, 2H), 7.85 (m, 2H), 8.05 (m, 2H)
RE 39	(4- <i>t</i> -Butylphenoxyethyl)-(4-chlorophenyl)-ketone	303	1.25 (m, 9H), 1.15 (s, 2H), 6.80 (m, 2H), 7.30 (m, 2H), 7.45 (m, 2H), 7.95 (M, 2H)
RE 40	(4-Phenylphenoxyethyl)-(phenyl)-ketone	289	5.25 (s, 2H), 7.00 (m, 2H), 7.20 – 7.25 (m, 1H), 7.40 (t, 2H), 7.50 – 7.60 (m, 6H), 7.60 (m, 1H), 8.00 (d, 2H)
RE 41	(Phenoxyethyl)-(4-phenylphenyl)-ketone	289	5.25 (s, 2H), 6.95 (m, 3H), 7.30 (M, 2H), 7.40 – 7.50 (m, 3H), 7.60 (m, 2H), 7.70 (m, 2H), 8.10 (d, 2H)
RE 42	(4-Chloro-2-carbamoylphenoxyethyl)-(4-fluorophenyl)-ketone	308	5.40 (s, 2H), 5.90 (bs, 1H), 6.90 (d, 1H), 7.20 (m, 2H), 7.40 (m, 1H), 8.00 (m, 2H), 8.30 (m, 1H), 8.70 (bs, 1H)
104	[2-( <i>N</i> -Phenylcarbamoyl)phenoxyethyl]-(4-fluorophenyl)-ketone	350	5.50 (s, 2H), 7.05 – 7.25 (m, 5H), 7.40 (m, 2H), 7.50 (m, 1H), 8.00 – 8.10 (m, 4H), 8.40 (d, 1H), 10.65 (bs, 1H)
105	[2-( <i>N</i> -Isopropylcarbamoyl)phenoxyethyl]-(4-fluorophenyl)-ketone	316	1.35 (d, 6H), 4.40 (quin, 1H), 5.40 (s, 2H), 6.95 (d, 1H), 7.10 – 7.30 (m, 3H), 7.40 (m, 1H), 8.00 – 8.01 (m, 2H), 8.30 (d, 1H), 8.70 (bs, 1H)

100804-2

- 87 -

106	[2-( <i>N</i> -Isobutylcarbamoyl)phenoxyethyl]-(4-fluorophenyl)-ketone	330	1.00 (d, 6H), 2.00 (quin, 1H), 3.40 (t, 2H), 5.40 (s, 2H), 6.95 (d, 1H), 7.10 – 7.30 (m, 3H), 7.45 (m, 1H), 8.00 (m, 2H), 8.30 (dd, 1H), 8.80 (bs, 1H)
107	(2,4-Dichloro-6-carbamoylphenoxyethyl)-(4-fluorophenyl)-ketone	344	5.40 (s, 2H), 5.80 (bs, 1H), 7.20 (m, 2H), 7.60 (m, 1H), 7.95 (m, 2H), 8.10 (m, 1H), 8.40 (bs, 1H)
108	[2-( <i>N,N</i> -Dimethylcarbamoyl)phenoxyethyl]-(4-fluorophenyl)-ketone	302	2.90 (s, 3H), 3.10 (s, 3H), 5.30 (s, 2H), 6.80 (d, 1H), 7.05 (t, 1H), 7.20 (m, 2H), 7.30 (d, 2H), 8.05 (m, 2H)
109	(2-Acetyl-amino-4-chlorophenoxyethyl)-(4-fluorophenyl)-ketone	322	2.25 (s, 3H), 5.35 (s, 2H), 6.90 (s, 1H), 7.00 (dd, 1H), 7.20 (m, 2H), 7.95 (m, 2H), 8.30 (d, 1H), 8.65 (bs, 1H)
110	(2-Carbamoylphenoxyethyl)-(4-chlorophenyl)-ketone	290	5.40 (s, 2H), 5.90 (bs, 1H), 6.95 (d, 1H), 7.15 (t, 1H), 7.45 – 7.60 (m, 3H), 7.90 (m, 2H), 8.30 (dd, 1H), 8.70 (bs, 1H)
111	(2,4-Dichloro-5-acetylaminophenoxyethyl)-(4-fluorophenyl)-ketone	356	2.20 (s, 2H), 5.35 (s, 2H), 7.10 – 7.20 (m, 2H), 7.40 (s, 1H), 7.55 (bs, 1H), 8.10 (m, 2H), 8.20 (bs, 1H)
112	(3-Acetylaminophenoxyethyl)-(4-chlorophenyl)-ketone	302	2.10 (s, 3H), 5.20 (s, 2H), 6.70 (d, 1H), 6.95 (d, 1H), 7.20 – 7.25 (m, 2H), 7.35 (bs, 1H), 7.45 (d, 2H), 7.95 (d, 2H)
113	(3-Carbamoylphenoxyethyl)-(4-chlorophenyl)-ketone	289	2.60 (s, 3H), 5.30 (s, 2H), 7.10 (m, 1H), 7.30 – 7.50 (m, 5H), 7.95 (d, 2H)
114	(3-Acetylaminophenoxyethyl)-(4-fluorophenyl)-ketone	286	2.10 (s, 3H), 5.20 (s, 2H), 6.70 (m, 1H), 6.95 (d, 1H), 7.10 – 7.20 (m, 3H), 7.30 (m, 2H), 8.05 (m, 2H)

100804-2

- 88 -

115	(3-Carbamoylphenoxyethyl)-(4-fluorophenyl)-ketone	272	5.30 (s, 2H), 7.10 – 7.20 (m, 3H), 7.40 (m, 2H), 7.45 (s, 1H), 8.05 (m, 2H)
116	(3-Acetylphenoxyethyl)-(4-fluorophenyl)-ketone		2.60 (s, 3H), 5.30 (s, 2H), 7.10 – 7.25 (m, 3H), 7.40 (t, 1H), 7.50 (m, 1H), 7.60 (m, 1H), 8.05 (m, 2H)
117	(3-Morpholinophenoxyethyl)-(4-fluorophenyl)-ketone	316	3.15 (t, 4H), 3.85 (t, 4H), 5.20 (s, 2H), 6.40 (m, 1H), 6.55 (m, 2H), 7.10 (m, 3H), 8.05 (m, 2H)
118	(2-Morpholinophenoxyethyl)-(4-fluorophenyl)-ketone	314	3.10 (t, 4H), 3.80 (t, 4H), 5.30 (s, 2H), 6.85 (m, 1H), 6.95 (m, 3H), 7.20 (m, 2H), 8.05 (m, 2H)
119	(4-Acetylaminoethoxyethyl)-(4-fluorophenyl)-ketone	288	2.15 (s, 3H), 5.20 (s, 2H), 6.85 (d, 2H), 7.00 – 7.20 (m, 2H), 7.40 (d, 2H), 8.05 (m, 2H)
120 <sup>2</sup>	(4-Chlorophenoxyethyl)-(3,5-difluorophenyl)-ketone	282 (M <sup>+</sup> )	
121 <sup>3</sup>	(2-Morpholinomethyl-3,5-dimethylphenoxyethyl)-(phenyl)-ketone	339 (M <sup>+</sup> )	
122 <sup>4</sup>	(2,4-Dibromophenoxyethyl)-(phenyl)-ketone	368 (M <sup>+</sup> )	
123 <sup>5</sup>	(2,4-Difluorophenoxyethyl)-(4-chlorophenyl)-ketone	282	
124 <sup>6</sup>	(2,4,6-Triiodophenoxyethyl)-(phenyl)-ketone	590 (M <sup>+</sup> )	
125 <sup>4</sup>	(2-Methoxy-4-propyl-5-bromophenoxyethyl)-(phenyl)-ketone	362 (M <sup>+</sup> )	

<sup>1</sup> MeCN instead of acetone was used as the solvent.<sup>2</sup> MeCN at room temperature rather than acetone at reflux was used as the solvent.<sup>3</sup> Using NaH in DMF as base.

100804-2

- 89 -

<sup>4</sup> Using KOH in ethanol.<sup>5</sup> DMF at room temperature as the solvent<sup>6</sup> KOH in n-butanol.**5 Reference Example 43****(4-Nitrophenoxyethyl)-(4-chlorophenyl)-ketone**

To a solution of 4-nitrophenol (42mg, 0.3mmol) and 4-chlorophenacyl bromide (84mg, 0.36mmol) in DCM was added MP-CO<sub>2</sub> resin solid resins (257mg, 0.9mmol). The reaction was shaken overnight. Scavenging reagents were added (131mg PS-thiophenol again  
 10 a resin, 34 mg MP-CO<sub>2</sub>) and the reaction was again shaken overnight. The reaction was filtered and the solvent was removed under reduced pressure. The resulting oil was purified by column chromatography (eluting with 10% EtOAc/isohexane to 50% EtOAc/isohexane) to yield an oil (35mg, 36%). NMR: 7.00 (d, 2H), 7.50 (d, 2H), 7.95 (d, 2H), 8.20 (d, 2H); m/z: 290 (M-H)<sup>+</sup>.

15

**Examples 126-152 and Reference Examples 44-50**

The procedure described in Reference Example 43 was repeated using the appropriate starting materials.

Ex	Compound	M/z
126	(4-Methoxyphenoxyethyl)-(4-chlorophenyl)-ketone	275 (M-H) <sup>+</sup>
RE 44	(2-Cyanophenoxyethyl)-(4-chlorophenyl)-ketone	270 (M-H) <sup>+</sup>
RE 45	(4-Chlorophenoxyethyl)-(4-chlorophenyl)-ketone	280 (M-H) <sup>+</sup>
127	(4-Chlorophenoxyethyl)-(2-methoxyphenyl)-ketone	275 (M-H) <sup>+</sup>
128	(4-Chlorophenoxyethyl)-(3-methoxyphenyl)-ketone	275 (M-H) <sup>+</sup>
RE 46	(4-Chlorophenoxyethyl)-(4-methoxyphenyl)-ketone	275 (M-H) <sup>+</sup>
RE 47	(4-Chlorophenoxyethyl)-(4-methylphenyl)-ketone	259 (M-H) <sup>+</sup>
RE 48	(4-Chlorophenoxyethyl)-(4-fluorophenyl)-ketone	263 (M-H) <sup>+</sup>
129	(4-Chlorophenoxyethyl)-(4-pentylphenyl)-ketone	317
130	(2-Fluorophenoxyethyl)-(4-chlorophenyl)-ketone	263 (M-H) <sup>+</sup>
RE 49	(2-Chlorophenoxyethyl)-(4-chlorophenyl)-ketone	280 (M-H) <sup>+</sup>
131	(2-Phenylphenoxyethyl)-(4-chlorophenyl)-ketone	321 (M-H) <sup>+</sup>
132	(2-Methylphenoxyethyl)-(4-chlorophenyl)-ketone	259 (M-H) <sup>+</sup>



100804-2

- 90 -

133	(2-Propylphenoxyethyl)-(4-chlorophenyl)-ketone	287 (M-H) <sup>+</sup>
134	(3-Cyanophenoxyethyl)-(4-chlorophenyl)-ketone	270 (M-H) <sup>+</sup>
135	(3-Methoxyphenoxyethyl)-(4-chlorophenyl)-ketone	275 (M-H) <sup>+</sup>
136	(3-Acetylphenoxyethyl)-(4-chlorophenyl)-ketone	287 (M-H) <sup>+</sup>
137	(3- <i>t</i> -Butylphenoxyethyl)-(4-chlorophenyl)-ketone	301 (M-H) <sup>+</sup>
138	(4-Acetylphenoxyethyl)-(4-chlorophenyl)-ketone	287 (M-H) <sup>+</sup>
139	(4-Cyclopentylphenoxyethyl)-(4-chlorophenyl)-ketone	313 (M-H) <sup>+</sup>
140	(3-Morpholinophenoxyethyl)-(4-chlorophenyl)-ketone	332 (M-H) <sup>+</sup>
141	(4-Butylphenoxyethyl)-(4-chlorophenyl)-ketone	303
142	(4-Morpholinophenoxyethyl)-(4-chlorophenyl)-ketone	332
143	[4-(2-methoxyethoxy)phenoxyethyl]-(4-chlorophenyl)-ketone	321
144	(4-Fluorophenoxyethyl)-(4-fluorophenyl)-ketone	249
145	(2-Isopropylphenoxyethyl)-(4-fluorophenyl)-ketone	273
146	(3-Chlorophenoxyethyl)-(4-fluorophenyl)-ketone	263 (M-H) <sup>+</sup>
147	[3-( <i>N,N</i> -Diethylamino)phenoxyethyl]-(4-fluorophenyl)-ketone	302
148	(3-Methoxyphenoxyethyl)-(4-fluorophenyl)-ketone	261
149	(4-Fluorophenoxyethyl)-(4-fluorophenyl)-ketone	247 (M-H) <sup>+</sup>
RE 50	(4-Methoxyphenoxyethyl)-(4-fluorophenyl)-ketone	259 (M-H) <sup>+</sup>
150	(4-Butylphenoxyethyl)-(4-fluorophenyl)-ketone	257 (M-H) <sup>+</sup>
151	(3-Isopropylphenoxyethyl)-(4-fluorophenyl)-ketone	273
152	(4-Morpholinophenoxyethyl)-(4-fluorophenyl)-ketone	316

**Example 153****(2-Hydroxyethylpyrid-5-yloxyethyl)-(phenyl)-ketone**

(*N*-Oxy-2-methylpyrid-5-yloxyethyl)-(phenyl)-ketone (Method 19; 4.72g, 5 19.4mmol) was dissolved in DMF (15ml) then chilled in an ice bath before the addition of trifluoroacetic anhydride (15ml). The solution was stirred at room temperature overnight. DCM (100ml) was added and reaction was carefully quenched with 2M sodium carbonate. The biphasic mixture was stirred at room temperature, and the resultant deep red solution was partitioned, the organic layer collected, dried (MgSO<sub>4</sub>), and the solvent was removed *in vacuo* 10 to give an oil. This was purified by column chromatography (EtOAc 100% to 2% methanol in EtOAc) to give the product (2.64g). NMR (DMSO-*d*<sub>6</sub>; 400MHz): 4.50 d, (2H), 5.25 (t, 1H),

100804-2

- 91 -

5.80 (s, 2H), 7.35 – 7.45 (m, 2H), 7.55 (m, 2H), 7.80 (t, 1H), 8.05 (dd, 2H), 8.25 (d, 1H); m/z 244.

### Reference Example 51

#### 5 (Phenoxyethyl)-(4-methylphenyl)-ketone

Sodium hydride (oil free, 1.85g, 77mmol) was added portionwise to phenol (6.6g, 70mmol) in DMF (100ml) at 0°C. When hydrogen evolution ceased,  $\alpha$ -bromo-4-methylacetophenone (14.9g, 70mmol) was added portionwise over 15 mins. The resulting dark solution was left to stand at 20°C for 4 hours. Water (500ml) was added and the mixture  
10 was extracted with EtOAc (2 x 200ml). The extracts were washed with water (200ml) and brine (200ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* to give an oil (16g). This was purified by column chromatography (5%EtOAc in hexane), and the resultant product recrystallized from hexane (5.44g, 34%). NMR: 2.40 (s, 3H), 5.2 (s, 2H), 6.9 – 7.0 (m, 2H), 7.2 – 7.35 (m, 5H), 7.9 (d, 2H); m/z 227.

15

### Reference Example 52

#### (1-Methylbenzimidazol-2-ylthiomethyl)-(4-bromophenyl)-ketone

(Benzimidazol-2-ylthiomethyl)-(4-bromophenyl)-ketone (Reference Example 53; 7g) was treated with sodium hydroxide (0.9g) in water (10ml). The reaction was stirred and  
20 methyl iodide was added (5ml). An insoluble solid formed which went into a sticky lump, and this was stirred overnight, during which it became a brownish colour. The solid was filtered under gravity, then stirred with DCM, then filtered under gravity again. The filtrate was discarded. The remaining solid was heated with ethanol (50ml), and filtered under gravity. The filtrate crystallised on standing overnight and the crystals were collected and dried in a  
25 warm cupboard. NMR (DMSO-d<sub>6</sub>): 3.75 (s, 3H), 5.08 (s, 2H), 7.1 – 7.6 (m, 6H), 7.8 – 8.2 (q, 2H).

### Reference Example 53

#### (Benzimidazol-2-ylthiomethyl)-(4-bromophenyl)-ketone

30 2-Mercaptobenzimidazole (3g) was suspended in acetone (100ml). 4-Bromophenacyl bromide was added and the whole amount set solid. A further 100ml of acetone was added and the reaction mixture swirled. The reaction mixture was filtered, to give a solid. Mp 241 – 245°C.

100804-2

- 92 -

**Example 154****(1-Methylimidazol-2-ylthiomethyl)-(4-chlorophenyl)-ketone**

4-Chloro- $\alpha$ -bromophenyl acetophenone (2.338g) and 1-methyl-2-mercaptoimidazole (1.14g) were dissolved in ethanol (50ml) and the reaction was stirred for 30 hours. The reaction was cooled in ice and quenched with 10% sodium acetate solution (80ml). The solid was filtered off to give 2 g product. This was recrystallized to give 140mg. Mp 68 - 70°C; NMR 3.6 (s, 3H), 4.50 (s, 2H), 6.9 - 8.0 (m, 7H).

**Example 155****(Pyrimidin-2-ylthiomethyl)-(4-bromophenyl)-ketone**

The procedure used above for Example 154 was repeated using 4-bromo- $\alpha$ -bromophenyl acetophenone and 2-mercaptopyrimidine to give the title compound. NMR (400MHz, DMSO- $d_6$ ): 5.00 (s, 2H), 7.40 (m, 1H), 8.00 (m, 2H), 8.20 (m, 2H), 8.80 (d, 2H); m/z 309.

**Example 156****{ $\alpha$ -[N-(Ethyl)-6-(bromo)naph-2-ylsulphonylamino]benzyl}-(phenyl)-ketone**

To [ $\alpha$ -(ethylamino)benzyl]-(phenyl)-ketone (Organic Reactivity (Tartu) (1984), 21(4), 418-27; 0.5mmol) was added DCM (3ml), followed by triethylamine (2mmol) in DCM (1ml). The mixture was vortexed for 10 seconds. 6-Bromonaph-2-ylsulphonylchloride (0.5mmol) in DCM (1ml) was then added. The reaction mixture was stirred at room temperature for 5 hours. The reaction mixture was washed with 2M hydrochloric acid (2 x 2ml) then water (2ml). The organic layer was separated in a Savant centrifugal evaporator to give the product (112.6mg). M/z 507 (M-H)<sup>+</sup>.

**Reference Example 54****(N-Methyl-4-chloroanilinomethyl)-(4-chlorophenyl)-ketone**

2-Bromo-4'-chloroacetophenone (233mg, 1.0mmol) was added to a solution of 4-chloro-N-methylaniline (295mg, 2.1mmol) in ethanol (8ml) and the mixture was stirred at room temperature overnight. The solid was filtered, washed with cold ethanol and dried to give the title compound as a solid (160mg, 0.55mmol). NMR : 3.1 (s, 3H), 4.7 (s, 2H), 6.6 (d, 2H), 7.1 (d, 2H), 7.4 (d, 2H), 7.9 (d, 2H); m/z 294.

100804-2

- 93 -

**Examples 157-160**

The procedure described in Reference Example 54 was repeated using the appropriate starting materials to obtain the compounds described below.

Ex	Compound	M/z	NMR
157	(N-Methyl-4-methylanilinomethyl)-(4-chlorophenyl)-ketone	274	
158	(N-Methoxy-4-methylanilinomethyl)-(4-chlorophenyl)-ketone	290	3.0 (s, 3H), 3.7 (s, 3H), 4.6 (s, 2H), 6.7 (d, 2H), 6.8 (d, 2H), 7.4 (d, 2H), 7.9 (d, 2H)
159	(N-Isopropyl-anilinomethyl)-(4-chlorophenyl)-ketone	288	1.2 (d, 6H), 4.2 (m, 1H), 4.6 (s, 2H), 6.6 (d, 2H), 6.7 (t, 1H), 7.1 (m, 2H), 7.4 (d, 2H), 8.0 (d, 2H)
160 <sup>1</sup>	(N-Methylanilinomethyl)-(4-iodophenyl)-ketone	352	

5 <sup>1</sup> Solvent used was dioxane not ethanol.

**Reference Example 55****[4-[1-(Pyrid-4-yl)piperazin-4-yl]phenethyl]-(4-methylphenyl)-ketone**

To a solution of 4-(4-pyridyl-1-piperazinyl)-benzaldehyde (WO 9728128; 1.0g, 3.75mmol) and 4-methylacetophenone (503mg, 3.75 mmol) in ethanol (25ml) was added concentrated aqueous sodium hydroxide solution (2 drops) and the mixture was stirred at room temperature overnight. The precipitated solid was filtered, washed with a small amount of cold ethanol, dried, taken up in ethanol (100ml) and hydrogenated over 10% Pd on charcoal. The catalyst was removed by filtration and the ethanol evaporated to leave a residue which was crystallised from BtOAc/hexane to give the title compound as a solid (320mg, 2.6mmol). Mp 114-115°C; C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O requires C; 77.9%; H;7.1%; N;10.9%; found C; 77.6%; H;7.1%; N;10.5%; m/z 386.

100804-2

- 94 -

**Example 161****(N-Mesyl-4-cyanoanilinomethyl)-(4-chlorophenyl)-ketone**

Sodium hydride (80mg of a 60% dispersion; 2.0mmol) was suspended in DMF (2ml) and treated with 4-methanesulphonamino-1-benzonitrile (400mg, 2.04mmol) at 0°C under argon. The mixture was stirred for 20 minutes then treated with a solution of 2-bromo-4'-chloroacetophenone (400mg, 1.72mmol) in DMF (2ml). The mixture was stirred at room temperature for 2 hours and then poured onto water (60ml). The aqueous layer was extracted with EtOAc (3x30ml), and the combined organic extracts washed with brine, dried and evaporated. The residue was purified by column chromatography using 30% EtOAc in hexane as eluent to give the title compound as a solid 500mg. NMR (300MHz): 3.2 (s, 3H), 5.2 (s, 2H), 7.5 (d, 2H), 7.55 (d, 2H), 7.6 (d, 2H), 7.90 (d, 2H); m/z 347 (M-H).

**Example 162****(Benzyl)-[4-(morpholinosulphonyl)phenyl]-ketone**

To 4-(morpholinosulphonyl)benzoylchloride (Method 20; 869mg-3mmol) in dimethoxyethane (10ml) was added bis(triphenylphosphine)-palladium chloride (211mg; 0.3mmol) and activated zinc (392mg; 6mmol). The mixture was degassed under argon and with stirring a solution of benzyl bromide (513mg; 3mmol) was added over 45 mins. After overnight at 20°C the mixture was diluted with EtOAc, washed with aqueous hydrochloric acid (2M, 25ml) and brine and dried (MgSO<sub>4</sub>). After concentration the residue was purified by chromatography with EtOAc-DCM as eluent to give a solid (524mg-51%). NMR: 3.00 (t, 4H), 3.75 (t, 4H), 4.3 (s, 2H), 7.3 (m, 5H), 7.8 (d, 2H), 8.15 (d, 2H).

**Example 163****25 [2-(Methoxymethylthio)benzoylaminomethyl]-(4-bromophenyl)-ketone**

2-(Methoxymethylthio)benzoic acid (Method 22; 376mg, 2mmol), dimethylaminopyridine (610mg, 5mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (400mg, 2.1mmol), 4-bromophenacylamine and hydrochloric acid (525mg, 2.1mmol) were dissolved in DMF (4ml). The reaction mixture was stirred overnight, evaporated and EtOAc was added (100ml). The EtOAc was washed with citric acid (3 x 50ml), saturated sodium hydrogencarbonate solution (50ml) and brine (50ml) and dried (MgSO<sub>4</sub>). The organic layer was evaporated to yield a yellow semi solid (670mg). This was

100804-2

- 95 -

purified by column chromatography (hexane: EtOAc 1:1) to give the title compound (219mg, 28%). M/z 396.

#### Example 164

##### 5 [N-(Thien-2-ylcarbonyl)anilinomethyl]-(4-fluorophenyl)-ketone

To a solution of (anilinomethyl)-(4-fluorophenyl)-ketone (Reference Example 56 230mg) in DCM (10ml) was added diisopropyl ethylamine (342 $\mu$ l) followed by 2-thiophene carbonyl chloride (106 $\mu$ l). The reaction was stirred at room temperature for 2 hours, then washed with hydrochloric acid (1M), saturated bicarbonate, water and brine. The organic  
10 layers were dried and evaporated. The solid was triturated with ether to give the title compound as a pale yellow solid (250mgs, 73%). M/z 340.

#### Reference Example 56

##### (Anilinomethyl)-(4-fluorophenyl)-ketone

15 To a suspension of aniline (910 $\mu$ l) and sodium bicarbonate (840mg) stirred at room temperature in ethanol (50ml) was added 4-fluorophenacyl bromide (2.17g). The reaction was stirred for 1 hour and the yellow suspension evaporated. The resultant slurry was taken up in water and extracted with EtOAc (2x50ml). The organics were dried, filtered and evaporated to give a yellow solid. This was triturated with isohexane to give a cream solid (1.45g, 63%).

20

#### Reference Example 57

##### (1,3-Diphenylprop-2-yl)-(phenyl)-ketone

To a stirred suspension of finely ground KOH in acetophenone was added Aliquat 336. The reaction was stirred for 5 mins at room temperature and then benzylbromide was  
25 added. The reaction was left to stir for 48 hours and then extracted with DCM. The DCM was washed with water and brine then dried (MgSO<sub>4</sub>), filtered and evaporated to yield a yellow oil. This material was first purified by column chromatography (50g silica, DCM) then prep HPLC to yield the product as an oil (107mg, 4%). NMR (DMSO-d<sub>6</sub>): 2.75 (m, 2H), 3.00 (m, 2H), 4.25 (m, 1H), 7.15 (m, 10H), 7.35 (t, 2H), 7.50 (m, 1H), 7.80 (d, 2H).

30

**Reference Example 58****(N-Methyl-4-methylphenylsulphonylaminomethyl)-(4-chlorophenyl)-ketone**

To a stirred solution of sodium hydride (60% suspension in mineral oil, 80mg, 2mmol) in anhydrous DMF (2ml) at 0°C under argon was added *N*-methyl-*p*-toluenesulphonamide (378mg, 2mmol). The reaction was stirred at 0°C for 20 mins and then a solution of 4-chlorophenacyl bromide (400mg, 1.7mmol) in DMF (2ml) was added. The reaction was allowed to warm to room temperature and left to stir for 2 hours. The reaction was quenched with cold water (60ml) and extracted with EtOAc (2x40ml), the combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and evaporated to give an oil. This oil was purified by column chromatography (eluting with 20% EtOAc/isohehexane) to yield the product as a solid (98mg, 17%). NMR(DMSO-d<sub>6</sub>): 2.40 (s, 3H), 2.70 (s, 2H), 4.70 (s, 2H), 7.40 (d, 2H), 7.60 (d, 2H), 7.70 (d, 2H), 8.00 (d, 2H); m/z: 338.

**Example 165****15 (N-Ethyl-4-methylphenylsulphonylaminomethyl)-(4-fluorophenyl)-ketone**

Using the procedure of Reference Example 58 using the appropriate starting materials the title compound was synthesised. NMR(DMSO-d<sub>6</sub>): 1.00 (t, 3H), 2.40 (s, 3H), 3.20 (q, 2H), 4.80 (s, 2H), 7.40 (m, 4H), 7.75 (d, 2H), 8.10 (m, 2H); m/z: 336.

**20 Preparation of Starting Materials**

The starting materials for the above Examples and Reference Examples are either commercially available or are readily prepared by standard methods from known materials. For example the following reactions are illustrations but not limitations of the preparation of some of the starting materials used in the above reactions.

25

**Method 1****N-Methoxy-N-methyl-3-thienylmethanamide**

Pyridine (5.0 ml) was added to a solution of N,O-dimethylhydroxylamine hydrochloride (3.00 g, 30.88 mmol) and 2-(3-thienyl)-acetyl chloride (3.55 g, 25.0 mmol) in DCM (100 ml) at 0°C. The resultant mixture was stirred at ambient temperature for 1 hour, washed with water (50 ml), dried and evaporated to dryness. The residue was purified by column chromatography using 30% EtOAc in hexane as eluent to give the title compound as a

100804-2

- 97 -

liquid (3.2 g, 17.3 mmol). NMR 3.2 (s, 3H), 3.6 (s, 3H), 3.8 (s, 2H), 7.0 (d, 1H), 7.1 (s, 1H), 7.2 (d, 1H).

#### Method 2

##### 5 N-Methoxy-N-methyl-2-thienylethanamide

Pyridine (0.5 ml) was added to a solution of N,O-dimethylhydroxylamine hydrochloride (300 mg, 3.08 mmol) and 3-(2-thienyl)-propionyl chloride (435 mg, 2.5 mmol) in DCM (10 ml) at 0°C. The resultant mixture was stirred at ambient temperature for 1 hour, washed with water (5 ml), dried and evaporated to dryness. The residue was purified by  
10 column chromatography using 30% EtOAc in hexane as eluent to give the title compound as a liquid (390 mg, 1.96 mmol); NMR: 2.8 (t, 2H), 3.2 (t, 2H), 3.2 (s, 3H), 3.6 (s, 3H), 6.8 (dd, 1H), 6.9 (dd, 1H), 7.1 (dd, 1H).

#### Method 3

##### 15 (3-Bromophenoxy)-(4-bromobenzyl)-ketone

3-Bromophenol (15g) and pyridine (12ml) were dissolved in DCM (75ml) and (4-bromophenyl)-acetyl chloride, dissolved in DCM (100ml), was added dropwise. The mixture was stirred overnight, then the reaction mixture was washed with water, 2M hydrochloric acid, 2M sodium hydroxide solution and brine. The reaction mixture was dried and the  
20 solvent removed *in vacuo*. The residue was purified by column chromatography (100% toluene) to give the required product (42g).

#### Method 4

##### N,N-Diethyl-N-( $\alpha$ -cyano-4-methoxybenzyl)amino

Para-anisaldehyde (2.76g, 20mmol) was dissolved in methanol and added over 1 hour to a solution of diethylamine hydrochloride (2.74g, 25mmol) and sodium cyanide (1.23g, 25mmol) in water (5ml). The solution was stirred at 30°C for 4 hours and then quenched with water (100ml) and extracted with ether. The extracts were washed with water, saturated sodium metabisulfite solution, water and brine. The solvent was removed *in vacuo* to give the  
30 product as a yellow oil (3.6g, 81%).



100804-2

- 98 -

**Method 5****(2-Fluoro-4-trifluoromethylphenyl)-(2-bromoprop-2-yl)-ketone**

(2-Fluoro-4-trifluoromethylphenyl)-(prop-2-yl)-ketone (Method 6) was placed in 48% hydrobromic acid (10ml), and bromine (1.8ml) was added dropwise over 6 hours. The reaction mixture was washed with water to remove excess bromine, and then passed through phase separation paper. The reaction mixture was then evaporated to dryness to give a crude yield of 10g. This product was used without further purification.

**Method 6****10 (2-Fluoro-4-trifluoromethylphenyl)-(prop-2-yl)-ketone**

1-(1-Hydroxy-2-methylpropyl)-2-fluoro-4-trifluoromethyl (Method 7; 10g, 42.7mmol) was dissolved in DCM (50ml). Pyridinium chlorochromate (13.8g, 64.1mmol) was added and the reaction was stirred at room temperature overnight. The reaction mixture was filtered through diatomaceous earth, triturating the remaining tar with DCM (3 x 30ml) and then evaporating to dryness. The resulting oil was dissolved in ether, filtering through diatomaceous earth, and evaporated to dryness to give a crude yield of 8g. The product was used without further purification.

**Method 7****20 1-(1-Hydroxy-2-methylpropyl)-2-fluoro-4-trifluoromethyl**

To *n*-butyl lithium, (28.5ml) in anhydrous ether (50ml) was added 4-bromo-3-fluorobenzotrifluoride (10g) in ether (50ml), dropwise at -70°C and under argon. The reaction was stirred for 15 mins, and a solution of butyraldehyde (2.95g) in ether (20ml) was added, and the reaction was stirred for a further 30mins. Acetic acid (10ml) in ether (20ml) was added while allowing the reaction mixture to warm to room temperature. Water (20ml) was added and the solution partitioned. The aqueous layer was washed with ether, the organic layers combined, dried and the solvent removed *in vacuo* to give a crude yield of 10g. The product was used without further purification.

30

100804-2

- 99 -

**Method 8****2-Chloro-5-acetylthiophene**

To a suspension of anhydrous aluminium trichloride (26.58g) in carbon tetrachloride (100ml) was added with vigorous stirring acetyl chloride (15.7g) over 20 mins with cooling in an ice bath. The resulting mixture was then treated with 2-chlorothiophene (23.7g), in carbon tetrachloride (25ml), with cooling in an ice bath over a 35 min period. The dark red solution was stirred a further 1 hour at 0°C and then poured onto ice/water/hydrochloric acid. The organic layer was separated and washed with water, dried and evaporated to give a mauve oil, which gradually solidified to a low melting solid (32g). This was used without further purification.

**Method 9****1-[1-(4-Chlorophenyl)-1-(trimethylsilyloxy)-1-(cyano)prop-2-yl]-1,2,4-triazole**

(4-Chlorophenyl)-[1-(1,2,4-triazol-1-yl)ethyl]-ketone (Method 10; 2.36g, 10mmol) was dissolved in toluene (15ml), under nitrogen. To this was added a catalytic amount of zinc iodide (200mg) and trimethylsilyl cyanide (1.2g, 1.6ml, 12mmol) and the reaction mixture stirred at room temperature for 2 hours. The mixture was heated 85°C and then left stirring at this temperature overnight. After cooling, ether was added to the reaction, and then the reaction mixture was washed with brine. The organic layer was separated, dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to gave an orange oil. The was purified by column chromatography to give the required product. M/z 334 (M<sup>+</sup>).

**Method 10****(4-Chlorophenyl)-[1-(1,2,4-triazol-1-yl)ethyl]-ketone**

Sodium hydride (62.5g, 50% dispersion, 2mol) was suspended in petroleum ether and washed. It was then freed from solvent using alternating vacuum and argon until dry and powder-like. Anhydrous DMF (200ml) was added under argon. A solution of triazole (90g, 2mol) in DMF (200ml) was added keeping the temperature between 20 and 30°C by cooling with an ice-bath. This was stirred for about 1.5 h until there was no further effervescence. A solution of (4-chlorophenyl)-(1-bromoethyl)-ketone (Method 11) in DMF (250ml) was added dropwise with stirring and cooling keeping the temperature between 20 and 30°C. The mixture was then stirred at room temperature overnight. The DMF was removed *in vacuo* and the resultant residue was portioned between ether and water. The ether extracts were washed

100804-2

- 100 -

with water, dried and the ether removed *in vacuo* to give a brown gum. A portion was triturated with petroleum ether to give a colourless solution which slowly deposited long needles. The rest of the material was purified by column chromatography (EtOAc) to give the product.

5

**Method 11****(4-Chlorophenyl)-(1-bromoethyl)-ketone**

4-Chloropropiophenone (122.5g) was dissolved in chloroform (500ml). Bromine (2ml) was added and the solution irradiated with a photoflood lamp until reaction began. This  
10 was cooled to about 10°C in an ice bath and the remainder of the bromine (35.5ml) was added dropwise, reacting almost immediately. After the addition, the ice bath was removed and the reaction was left to stand overnight. The reaction mixture was washed with water (2 x 200ml), saturated sodium hydrogen carbonate and water. The organic layer was then dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to give the crude product. This was recrystallized from  
15 cyclohexane to give the product as white crystals.

**Method 12****N-(Isopropyl)-N-(mesyl)cyclohexylamino**

To a stirred solution of N-isopropylcyclohexylamine (2g, 0.014mol) and triethylamine  
20 (1.49g, 0.015mol) in DCM (80ml) at 0°C was added mesylchloride (1.62g, 0.014mol). The reaction was stirred at 0°C for 10 minutes then allowed to warm to room temperature and left to stir for a further 30 minutes. The reaction mixture was transferred to a separating funnel and diluted to ~150ml with DCM. The solution was then washed with hydrochloric acid (2M; 50ml), water (50ml) and brine (30ml), dried (MgSO<sub>4</sub>), filtered and evaporated to yield the  
25 product as a clear oil (2.26g, 74%). NMR: 1.05 (br m, 1H), 1.30 (br d, 8H), 1.60 (br s, 2H), 1.80 (br s, 6H), 2.85 (s, 3H), 3.30 (br m, 1H), 3.80 (m, 1H); m/z: 219.

**Method 13****N-(Methyl)-N-(mesyl)pyrid-2-ylamino**

30 To a stirred solution of 2-(methylamino)pyridine (2g, 0.018mol) and triethylamine (1.82g, 0.018mol) in anhydrous DCM (80ml) at 0°C was slowly added mesyl chloride (2.06g, 0.018mol). The reaction was stirred at 0°C for 10 mins then allowed to warm to room temperature and stirred for a further 30 mins. The solvent was removed under reduced

100804-2

- 101 -

pressure and resulting solid was partitioned between ether and water. The organic layer was separated and reextracted with ether. The combined organic layers were washed with brine then dried (MgSO<sub>4</sub>), filtered and evaporated to yield an oil (2.2g, 67%). NMR: 3.00 (s, 3H), 3.40 (s, 3H), 7.15 (br m, 1H), 7.45 (br m, 1H), 7.75 (br m, 1H), 8.45 (br s, 1H).

5

**Method 14****1-(Methyl)-4-(mesyl)piperazine**

To a stirred solution of 1-methylpiperazine (1g, 10mmol) and triethylamine (1.11g, 11mmol) in anhydrous DCM (70ml) at 0°C was added mesyl chloride (1.15g, 10mmol). The reaction was stirred at 0°C for ten minutes then allowed to warm to room temperature and stirred for a further 30 minutes. The volatiles were removed under reduced pressure and the resulting material was partitioned between DCM and 2M NaOH. The organic layer was separated and washed with brine, dried (MgSO<sub>4</sub>), filtered and evaporated to yield an oil (962mg, 53%). NMR: 2.35 (s, 3H), 2.50 (t, 4H), 2.75 (s, 3H), 3.30 (m, 4H).

15

**Method 15****N-(Methyl)-N-(mesyl)-4-chloroaniline**

To a stirred solution of 4-chloro-N-methylaniline (705mg, 5mmol) in anhydrous pyridine (4ml) at 0°C was added dropwise methanesulphonyl chloride (0.41ml, 5.25mmol). The reaction was allowed to warm to room temperature and was stirred for 1 hour. The reaction mixture was partitioned between ether (25ml) and 2M hydrochloric acid (30ml). The aqueous layer was extracted with ether (2x25ml). The combined organic extracts were washed with 1M hydrochloric acid, saturated sodium bicarbonate and brine, dried (MgSO<sub>4</sub>), filtered and evaporated. The resulting material was crystallised from EtOAc/iso hexane to yield a white solid (900mg, 83%). NMR: 2.85 (s, 3H), 3.30 (s, 3H), 7.35 (m, 4H).

25

**Methods 16 - 18**

The procedure described in Method 15 was repeated using the appropriate anilines to replace the 4-chloro-N-methylaniline to obtain the compounds described below.

30

100804-2

- 102 -

Ex	Compound	NMR
16	<i>N</i> -(Methyl)- <i>N</i> -(mesyl)-4-methoxyaniline	2.80 (s, 3H), 3.30 (s, 3H), 3.80 (s, 3H), 6.90 (d, 2H), 7.30 (d, 2H)
17	<i>N</i> -(Ethyl)- <i>N</i> -(mesyl)-4-methoxyaniline	1.10 (t, 3H), 2.85 (s, 3H), 2.65 (q, 2H), 3.80 (s, 3H), 6.90 (d, 2H), 7.25 (d, 2H)
18	<i>N</i> -(mesyl)-4-chloroaniline	3.00 (s, 3H), 6.70 (br s, 1H), 7.20 (d, 2H), 7.35 (d, 2H)

**Method 19****(*N*-Oxy-2-methylpyrid-5-yloxymethyl)-(phenyl)-ketone**

- 5 (2-Methylpyrid-5-yloxymethyl)-(phenyl)-ketone (Example 101; 5.7g, 25.1mmol) was stirred in DCM (50ml) then chilled in an ice bath before adding meta-chloroperoxybenzoic acid (9.5g, 27.6mmol @ 50%) and the reaction was stirred at room temperature overnight. A thick precipitate formed, this was filtered off and washed with ether to give the title compound. The filtrate was evaporated to give a solid which was washed with ether to give a
- 10 second crop of the product (5.22g).

**Method 20****4-(Morpholinosulphonyl)benzoylchloride**

- 4-(Morpholinosulphonyl)benzoic acid (Method 21; 3.6g; 0.0133mol) was heated at
- 15 reflux for 10 hours in thionyl chloride (50ml) containing 1drop of DMF. After concentration the title compound was obtained as a solid ( 3.54g-92%) which was used without purification.

**Method 21****4-(Morpholinosulphonyl)benzoic acid**

- 20 To a suspension of 4-carboxyphenylsulphonylchloride (2.2g; 0.01mol) in DCM (25ml) was slowly added morpholine (4.35ml; 0.05mol) at gentle reflux. After 2 hours at room temperature the DCM was evaporated off. The residue was acidified with excess of aqueous hydrochloric acid. The solid was filtered, washed with water and dried under vacuum over P<sub>2</sub>O<sub>5</sub>. A solid was obtained (2.2g-81%). NMR (DMSO-d<sub>6</sub>) 2.9 (t, 4H), 3.6 (t, 4H), 7.8 (d,
- 25 2H), 8.2 (d, 2H).

100804-2

- 103 -

**Method 22****2-(Methoxymethylthio)benzoic acid**

To a stirred solution of powdered potassium hydroxide (1.68g, 30mmol), triethylbenzylammonium bromide (0.23g, 10mol%) in dry methanol (40ml) and 5 bromochloromethane (40ml) was added a solution of thiosalicic acid (1.54g, 10mmol) in dry methanol (40ml). The solution was stirred overnight at room temperature under an inert atmosphere. Hydrochloric acid (1M; 75ml) was added and the organics were extracted with DCM: ether (1:1). The organics were pooled and washed with brine (50ml) dried (MgSO<sub>4</sub>) and evaporated to yield a pale yellow solid (1.85g, 93%). M/z 197.

10

**Method 23****1-(4-Fluorophenyl)-2-methoxyethanone**

To a 1.0M solution of 4-fluorophenylmagnesium bromide in THF (24.0 ml, 24.0 mmol) at 0°C was added a solution of methoxyacetonitrile (1.42g, 20.0 mmol) in ether (15 15 ml). The resultant mixture was stirred at ambient temperature for 2 hours and then quenched with 1M aqueous hydrochloric acid (40 ml). This mixture was stirred at ambient temperature for 2 hours and the aqueous layer was extracted with ether (2x40 ml). The combined extracts were washed with saturated aqueous sodium hydrogen carbonate solution, brine, dried and evaporated. The residue was purified by column chromatography using 10% EtOAc in hexane 20 as eluent to give the title compound as a solid (2.0 g, 11.9 mmol). NMR 3.5 (s, 3H), 4.7 (s, 2H), 7.1 (m, 2H), 8.0 (m, 2H).

**Method 24****[1-(2,4-Dichlorophenyl)vinyl]-(4-chlorophenyl)-ketone**

25 (2,4-Dichlorobenzyl)-(4-chlorophenyl)-ketone (Reference Example 17; 15g, 50mmol) was stirred in tetramethyldiaminomethane (25ml) and cooled in an ice-bath. Acetic anhydride (25ml) was added dropwise, keeping the temperature below 40°C. When the addition was complete, the ice bath was removed and the reaction stirred at room temperature for 1 hour. The reaction mixture was poured slowly onto crushed ice in water (500ml) with stirring. The 30 product precipitated and this was collected by filtration and dried in a dessicator. Mp 90 - 93°C; NMR: 6.00 (d, 2H), 7.35 (m, 3H), 7.75 (q, 4H).

- 104 -

**Method 25****[2-(4-Fluorophenyl)vinyl]-(4-trifluoromethylphenyl)-ketone**

Sodium hydroxide (1g, 25mmol) was dissolved in a mixture of water (80ml) and ethanol (20ml). 4-Trifluoromethylacetophenone (3.6g, 20mmol) was added followed by the rapid addition of 4-fluorobenzaldehyde (2.14ml, 2.48g, 20mmol) during which the temperature of the reaction was maintained at about room temperature with a water bath. The reaction was stirred at room temperature for 2 hours during which time an oily solid precipitated. The reaction vessel was stored in the fridge overnight. The product was filtered off, washed with water until the pH was neutral and then recrystallized from ethanol with a few drops of water to give a yellowish solid (3.25g, 55%). Mp 89 - 92°C.

**Methods 26-29**

The procedure described in Method 25 was carried out using the appropriate starting materials to obtain the products described below.

Meth	Compound	Data
26	[2-(4-Fluorophenyl)vinyl]-(4-chlorophenyl)-ketone	Mp 135°C
27	[2-(4-Chlorophenyl)vinyl]-(2,4-difluorophenyl)-ketone	Mp 102°C
28	[2-(4-Fluorophenyl)vinyl]-(2,4-difluorophenyl)-ketone	Mp 75 - 76°C
29 <sup>1</sup>	[2-(4-Methoxyphenyl)vinyl]-(phenyl)-ketone	


<sup>1</sup> This compound was prepared using lithium hydroxide as the base.

100804-2

- 105 -

Claims

1. The use of a compound of formula (I):



5

wherein:

Ring A is selected from aryl or heteroaryl;

- $R^1$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkanoyl,  $C_{1-6}$ alkanoyloxy,  $N-(C_{1-6}alkyl)amino$ ,  $N,N-(C_{1-6}alkyl)_2amino$ ,  $C_{1-6}alkanoylamino$ ,  $N-(C_{1-6}alkyl)carbamoyl$ ,  $N,N-(C_{1-6}alkyl)_2carbamoyl$ ,  $C_{1-6}alkylS(O)_a$ , wherein a is 0 to 2,  $C_{1-6}alkoxycarbonyl$ ,  $N-(C_{1-6}alkyl)sulphamoyl$ ,  $N,N-(C_{1-6}alkyl)_2sulphamoyl$ ,  $C_{1-6}alkylsulphonylamino$ , carbocyclyl, heterocyclyl, carbocyclyl $C_{0-6}alkylene-Y-$  and heterocyclyl $C_{0-6}alkylene-Y-$ ; or two  $R^1$  on adjacent carbons may form an oxy $C_{1-4}alkoxy$  group; wherein  $R^1$  may be optionally substituted on carbon by one or more groups selected from  $R^7$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^8$ ;

- 15  $n$  is 0-3; wherein the values of  $R^1$  may be the same or different;

- $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  are independently selected from hydrogen, hydroxy, amino, cyano,  $C_{1-4}alkyl$ ,  $C_{1-4}alkoxy$ ,  $N-(C_{1-4}alkyl)amino$ ,  $N,N-(C_{1-4}alkyl)_2amino$ ,  $C_{1-4}alkylS(O)_a$ , wherein a is 0 to 2,  $C_{1-4}alkoxycarbonyl$ , carbocyclyl, heterocyclyl, carbocyclyl $C_{1-4}alkyl$  and heterocyclyl $C_{1-4}alkyl$ ; or  $R^2$  and  $R^3$  together form oxo; wherein  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  may be independently optionally substituted on carbon by one or more groups selected from  $R^9$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{10}$ ;

- 25  $X$  is  $-CR^{11}R^{12}$ -,  $-S(O)_a$ -,  $-O$ -,  $-NR^{13}$ -,  $-C(O)$ -,  $-C(O)NR^{14}$ -,  $-NR^{15}C(O)$ -,  $-SO_2NR^{16}$ - or  $-NR^{16}SO_2$ -; wherein a is 0 to 2;

$r$  is 1 or 2;

$q$  is 0 or 1;

$p$  is 0 or 1;



100804-2

- 106 -

**Ring B** is carbocyclyl or heterocyclyl; wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{17}$ ;

$R^6$  is a substituent on carbon and is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl, heterocyclyl, carbocyclylC<sub>0-4</sub>alkylene-Y- and heterocyclylC<sub>0-4</sub>alkylene-Y-; wherein  $R^6$  may  
10 be optionally substituted on carbon by one or more groups selected from  $R^{18}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{19}$ ;

m is 0-3; wherein the values of  $R^6$  may be the same or different;

Y is -S(O)<sub>a</sub>-, -O-, -NR<sup>20</sup>-, -C(O)-, -C(O)NR<sup>21</sup>-, -NR<sup>22</sup>C(O)- or -SO<sub>2</sub>NR<sup>23</sup>-; wherein a is  
15 0 to 2;

$R^7$ ,  $R^9$  and  $R^{18}$  are independently selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  
20  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub> wherein a is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-4}$ alkylsulphonylamino, carbocyclyl and heterocyclyl; wherein  $R^7$ ,  $R^9$  and  $R^{18}$  may be independently optionally substituted on carbon by one or more  $R^{26}$ ;

$R^{11}$  and  $R^{12}$  are independently selected from hydrogen, hydroxy, amino, cyano,  
25  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino, carbocyclyl, heterocyclyl carbocyclylC<sub>1-4</sub>alkyl, heterocyclylC<sub>1-4</sub>alkyl; wherein  $R^{11}$  and  $R^{12}$  may be independently optionally substituted on carbon by one or more groups selected from  $R^{24}$ ; and wherein if said heterocyclyl contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from  $R^{25}$ ;

30  $R^{24}$  is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy,  $C_{1-4}$ alkyl,  $C_{2-4}$ alkenyl,  $C_{2-4}$ alkynyl,  $C_{1-4}$ alkoxy,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkanoyloxy,  $N$ -( $C_{1-4}$ alkyl)amino,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>amino,  $C_{1-4}$ alkanoylamino,  $N$ -( $C_{1-4}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-4}$ alkylS(O)<sub>a</sub>

100804-2

- 107 -

wherein  $a$  is 0 to 2,  $C_{1-4}$ alkoxycarbonyl,  $N$ -( $C_{1-4}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-4}$ alkyl)<sub>2</sub>sulphamoyl and  $C_{1-4}$ alkylsulphonylamino;

$R^8$ ,  $R^{10}$ ,  $R^{17}$ ,  $R^{19}$  and  $R^{25}$  are independently selected from  $C_{1-4}$ alkyl,  $C_{1-4}$ alkanoyl,  $C_{1-4}$ alkylsulphonyl,  $C_{1-4}$ alkoxycarbonyl, carbamoyl,  $N$ -( $C_{1-4}$ alkyl)carbamoyl, 5  $N,N$ -( $C_{1-4}$ alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl, heterocyclyl and phenylsulphonyl;

$R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$  and  $R^{23}$  are independently selected from hydrogen, phenyl,  $C_{1-4}$ alkylsulphonyl and  $C_{1-4}$ alkyl;

$R^{26}$  is selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, 10 amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxymethyl, methylamino, ethylamino, dimethylamino, diethylamino,  $N$ -methyl- $N$ -ethylamino, acetylamino,  $N$ -methylcarbamoyl,  $N$ -ethylcarbamoyl,  $N,N$ -dimethylcarbamoyl,  $N,N$ -diethylcarbamoyl,  $N$ -methyl- $N$ -ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, 15  $N$ -methylsulphamoyl,  $N$ -ethylsulphamoyl,  $N,N$ -dimethylsulphamoyl,  $N,N$ -diethylsulphamoyl or  $N$ -methyl- $N$ -ethylsulphamoyl; or a pharmaceutically acceptable salt thereof; in the manufacture of a medicament for use in the inhibition of 11 $\beta$ HSD1; with the proviso that said compound is not (1-methyl-1-pyrid-3-ylethyl)-(pyrid-3-yl)-ketone.


20

100804-2

- 108 -

**ABSTRACT****TITLE: CHEMICAL COMPOUNDS**

5      Compounds of formula (I):



(I)

wherein variable groups are as defined within; for use in the inhibition of 11 $\beta$ HSD1 are described.

10

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**